

Appendix 2. Plants on and in the vicinity of the Bering Land Bridge during the Late Cretaceous and Tertiary. References: 1= Budantsev, 1997; 2= Penhallow, 1908; 3= LaMotte, 1952; 4= Knowlton, 1910; 5= Heer, 1855; 6= Chaney, 1951; 7= Lesquereux, 1874; 8= Berry, 1926; 9= Knowlton, 1899; 10= Wolfe and Wehr, 1987; 11= Heer, 1876; 12= Lesquereux, 1878; 13= Boyd, 1990; 14= Manchester et al., 2009; 15= Herman, 1990; 16= Golovneva, 2000 (Cretaceous-Paleogene boundary); 17= Stockey et al., 1997; 18= Spicer et al., 1994; 19= Shezepetov and Golovneva, 2010; 20= Herman and Spicer, 2010 (Middle Cretaceous); 21= Budantsev, 1994 [Paleogene (Paleocene-Eocene)]; 22= Lavrenko and Fot'janova 1994; 23= Lebedev, Ye. L., 1992; 24= Vakhrameev, 1991; 25= Zhilin (ed.), 1991 (see Appendix note 1); 26= Velichko et al., 2005; 27= Arkhipov et al., 2005; 28= Fradkina et al., 2005a; Fradkina et al., 2005b; 29= Akhmetiev, 2010 (see also Akhmetiev, 2007); 30= Pavlyutkin and Chekryzhov, 2007; 31= Herman, 2013 (all floras and taxa are listed here under K although some may extend into the Pa (Danian)? Fide Herman, 2013; Herman et al., 2004); 32= Hollick, 1930; 33= Bell, 1949; 34= Brink et al., 2009; 35= Wolfe et al., 1966; 36= Stockey et al., 2001; 37= Wolfe, 1977; 38= Vakhrameev, 1958; 39= Pigg et al., 2001, 2007; 40= Harris et al., 2014; 41= Piel, 1971; 42= Wolfe and Tanai, 1987; 43= Becker, 1969; 44= Leopold and Liu, 1994 (Miocene and probably early Pliocene); 45= Wolfe and Tanai, 1980; 46= Reinink-Smith and Leopold, 2005; 47= Chelebaeva, 1978; 48= Pound et al., 2015; 49= Spicer and Herman, 2001; 50= Spicer and Herman, 1996; 51= Moiseeva, et al., 2009; 52= Takhtajan and others (1974 et seq.); 53= McIver and Basinger, 1993 (Paleocene, Saskatchewan, Canada); 54= Bell, 1949 (Cretaceous/Paleocene; see pp. 17, 29 for age assignments of formations), western Alberta, Canada; 55= Falcon-Lang et al., 2016. Where the LaMotte (3) or Takhtajan and others (52) catalogs are cited as a source of the listing for a genus these are indicated in brackets ([-]). Numbers/letters immediately after

plant names refer to Notes at the end of the Appendices. Numbers in the right columns refer to the above references. w= widespread. Taxa are as listed by the original author. General catalogs of fossil plants are Lamotte (1952) for North America; Takhtajan (1974, 1982) and Budantsev (1994, 2005) for Russia; and Graham (2010, p. xiii; 2012, URL: <http://www.mobot.org/mobot/research/CatalogFossil/catalog.shtml>) for the New World and references therein to global databases. Note that three genera listed in Doyle and Hickey (1976) for the Potomac (mid-Cretaceous) Group of the Atlantic Coastal Plain also occur in Beringia (*Araliaephylum*, *Protophyllum*, *Sapindopsis*). The complete inventory for the Potomac Group is included in Appendix 3 with the North Atlantic Land Bridge. Taxa listed in this and other Appendices have been extracted from more complete lists (I, II). Most of those identified by an artificial nomenclature with only mention of presumed biological affinities are not included.

Taxon	West Beringia (Siberia/Kamchatka)	East Beringia (Western North America)
<u>Cretaceous (Late)</u>		

Ferns and allied groups

<i>Acrostichopteris</i> aff. <i>parvifolia</i>	x (31)
<i>A. pluripartita</i>	x (31)
<i>Adiantopteris gracilis</i>	x (31)
<i>Adiatum? paululum</i>	x (54)
<i>Anemia arctopteroides</i>	x (24)

<i>A. cf. longata</i>	x (31)
<i>Arctopteris heteropinnula</i>	x (31)
<i>A. cf. kolymensis</i>	x (20, 31, 49) x (31, 49)
<i>A. lenaensis</i>	x (31)
<i>A. obtusipinnatus</i>	x (31)
<i>A. penzhinensis</i>	x (31) x (31)
<i>A. rarinervis</i>	x (31, 49) x (31, 49)
<i>A. aff. rarytkinensis</i>	x (31)
<i>Asplenium dicksonianum</i>	x (20, 24, 31) x (31)
<i>A. forester</i>	x (32)
<i>A. popovii</i>	x (31)
<i>A. rigidum</i>	x (20, 31)
<i>Azimia</i>	x (24)
<i>Birisia acutana</i>	x (31) x (31)
<i>B. alata</i>	x (31, 49) x (31, 49)
<i>B. jelisejevii</i>	x (31) x (31)
<i>B. ochotica</i>	x (31, 49) x (31, 49)
<i>B.(?) oerstedtii</i>	x (31) x (31)
<i>B. onychioides</i>	x (20, 24, 31) x (31)
<i>Cladophlebis acuta</i>	x (31)
<i>C. argutula</i>	x (31)
<i>C. columbianus</i>	x (31)
<i>C. contrarius</i>	x (31)

<i>C. cf. frigida</i>	x (31)	x (31)
<i>C. gluschinskii</i>	x (31)	
<i>C. aff. grandis</i>	x (31)	
<i>C. groenlandica</i>		x (33)
<i>C. haiburnensis</i>	x (31)	
<i>C. cf. kolymensis</i>		x (31)
<i>C. cf. lobulata</i>	x (31)	x (31)
<i>C. meyerii</i>	x (20)	
<i>C. pseudolobifolia</i>	x (31)	
<i>C. septentrionalis</i>	x (31)	x (31, 32)
<i>C. speranskae</i>	x (31)	
<i>C. ubiencaensis</i>	x (31)	
<i>C. venustus</i>	x (31)	
<i>C. virginensis</i>		x (31)
<i>C. aff. williamsonii</i>	x (31)	
<i>Cladophlebidium singulinerve</i>	x (31)	
<i>Coniopteridium</i>	x (31)	
<i>Coniopteris anadyrensis</i>	x (31)	
<i>C. arctica</i>	x (31)	x (31)
<i>C. aff. bicrenata</i>	x (31)	x (31)
<i>C. cf. brevifolia</i>	x (31)	x (31)
<i>C. cf. C. compressa</i>	x (20, 31)	x (31)
<i>C. dicksonioides</i>	x (31)	

<i>C. gracillima</i>	x (31)	
<i>C. (Birisia?) grebенcaensis</i>	x (31)	x (31)
<i>C. aff. ketovae</i>	x (31)	
<i>C. aff. maakiana</i>	x (31)	x (31)
<i>C. cf. nympharum</i>	x (20, 31)	
<i>C. cf. saportana</i>	x (20, 31)	x (31)
<i>C. setacea</i>		x (31)
<i>C. tschuktschorum</i>	x (31)	
<i>C. yukonensis</i>		x (31)
<i>Danaeites kinesis</i>	x (24)	
<i>Dennstaedtia blomstrandii</i>		x (33)
<i>D. sp.</i>	x (31)	
<i>Elatocladus</i>		x (31)
<i>Equisetites burejensis</i>	x (31, 49)	x (31, 49)
<i>E. ramosus</i>	x (31)	
<i>Equisetum arcticum</i>	x (31)	x (31, 33)
<i>E. perlaevigatum</i>		x (54)
<i>Filicites knowltoni</i>		x (54)
<i>Gleichenia crenata</i>		x (31)
<i>G. pseudocrenata</i>	x (31, 49)	x (31, 49)
<i>Gleichenites asiatica</i>	x (31)	
<i>G. gieseckianus</i>		x (31)
<i>G. microphylla</i>	x (31)	x (31)

<i>G. onkilonica</i>	x (31)	
<i>G. zippei</i>	x (31)	x (31)
<i>Goniopteris arctica</i>	x (49)	x (49)
<i>Hausmannia</i> aff. <i>atwoodii</i>	x (31)	[3]
<i>H. bilobata</i>	x (31)	
<i>H. bipartita</i>	x (31)	
<i>H. nansenii</i>	x (20)	
<i>H. cf. undulata</i>	x (31)	
<i>Isöetites</i>	x (16)	[3]
<i>Kolymella</i> aff. <i>raevskii</i>		x (31)
<i>Lycopodium</i>		x (31)
<i>Ochotopteris ochotensis</i>		x (31)
<i>O. kamtchatica</i>	x (31)	
<i>Onoclea hesperia</i>	x (31)	x (31, 33)
<i>Onychiopsis elongata</i>	x (15, 31)	x (31)
<i>O. grandipinnula</i>	x (31)	
<i>O. psilotoides</i>	x (31, 49)	x (31, 49)
<i>Osmunda</i> cf. <i>cretacea</i>	x (31)	x (31)
<i>O. cf. denticulata</i>	x (31)	x (31)
<i>O. macrophylla</i>		x (33)
<i>O. serrulata</i>	x (31)	
<i>Osmundopsis efimovae</i>	x (31)	
<i>Penzhinopteris pergamentii</i>	x (31)	

<i>Ruffordia goeppertii</i>	x (31)	x (31)
<i>R. magnifolia</i>	x (31)	
<i>Salvinia</i>	x (24)	[3]
<i>Schizaea</i>	x (31)	
<i>Scleropteris verchojanensis</i>	x (31)	
<i>Selaginella</i>	x (31)	[3]
<i>Sphenopteris crispans</i>	x (31)	
<i>S. cf. dentata</i>		x (31)
<i>S. mclearnii</i>	x (31)	
<i>Tchaunia filatovii</i>		x (31)

Gymnosperms

<i>Abies</i> (10)	x (24)	[3]
<i>Abietites</i>		x (31)
<i>Agathis tollii</i>	x (20)	
<i>Alatispermum malandinii</i>	x (20)	
<i>Anomozamites arcticus</i>	x (20, 31)	
<i>A. sp.</i>	x (24)	x (31)
<i>Araucarites anadyrensis</i>	x (15, 23, 31)	x (31)
<i>A. conophyllus</i>	x (31)	
cf. <i>Athrotaxites berryi</i>	x (31)	
<i>Athrotaxopsis expansa</i>	x (15, 31)	x (31)
<i>A. grandis</i>	x (31)	x (31)

<i>Baiera ahnertii</i>	x (31)	
<i>B. cf. gracilis</i>	x (31)	x (31)
<i>Brachiphyllum</i>	x (20, 24)	
<i>Butefia obliqua</i>	x (31)	x (31)
<i>Cedrus lopatinii</i>	x (20)	[3]
<i>C. sp.</i>	x (24, 31)	
<i>Cephalotaxopsis borealis</i>	x (31)	
<i>C. heterophylla</i>	x (20, 31)	
<i>C. intermedia</i>	x (24, 31, 49)	x (31, 32, 49)
<i>C. microphylla</i>		x (31, 32, 33)
<i>C. sangarensis</i>	x (31)	
<i>Cephalotaxopsis</i> (= <i>Taxites</i> fide Vakhrameev, 1991, p. 182)	x (15, 24)	
<i>Cryptomeria subulata</i>	x (31)	x (31)
<i>Cryptomerites ancistrophyllus</i>	x (31)	
<i>C. lambii</i>		x (33)
<i>Ctenis</i>	x (20, 24, 49)	x (31, 49)
<i>Cunninghamia hornbyensis</i>		x (34)
<i>C. sp.</i>	x (24)	
<i>Cupressinocladus cretaceous</i>	x (31)	x (31)
[genus as <i>Thuja</i> by Bell (33) fide Vakhrameev (24) p. 212]		x (33)
<i>Cycadites hyperborea</i>	x (31)	

<i>Cyparassisium</i>	x (24)	
<i>Czekanowskia rigida</i>	x (20?, 23, 31)	x (31)
<i>Desmiophyllum</i> (Gymnospermae, incertae sedis) <i>D. magnum</i>	x (31)	x (31)
<i>D. rigidum</i>	x (20)	
<i>Drepanolepsis</i>	x (31)	
<i>Elatides curvifolia</i>		x (31)
<i>Elatocladus</i> aff. <i>manchurica</i>	x (20, 31)	x (31)
<i>E. gracillimus</i>	x (31)	
<i>E. intermedius</i>		x (54)
<i>E. olriki</i>		x (54)
<i>E. smittiana</i>	x (15, 31)	x (31)
<i>E. talensis</i>	x (31)	
<i>E. ungeri</i>		x (54)
<i>Encephalartopsis vassilevskajae</i>	x (31)	
<i>Florinia borealis</i>	x (20, 31)	x (31)
<i>Fokieniopsis</i> aff. <i>catenulata</i>		x (31)
<i>Ginkgo adiantoides</i>	x (20, 24, 31, 49)	x (31, 49)
<i>G. angusticuneata</i>	x (31)	
<i>G. concinna</i>	x (31, 49)	x (31, 49)
<i>G. delicata</i>	x (31)	
<i>G. huttoni</i>	x (31)	x (31)
<i>G. lepida</i>	x (31)	x (31)

<i>G. minor</i>		x (32)
<i>G. nana</i>	x (31)	
<i>G. obrutschewii</i>	x (20	
<i>G. paradiantoides</i>	x (31)	
<i>G. pilifera</i>	x (2)	
<i>G. polaris</i>	x (31)	
<i>G. sibirica</i>	x (31)	x (31)
<i>G. singularis</i>	x (31)	
<i>Ginkgoites adiantoides</i>		x (33,54)
<i>G. sibirica, G. volnovii</i>	x (20)	
<i>Glyptostrobus groenlandicus</i>	x (31)	x (31)
<i>G. nordenskioldii</i>	x (31)	
<i>Heilungia gaimanensis</i> (now includes <i>Butefia</i> , see ref. 19)	x (19)	
<i>H. cf. oloensis</i>	x (49)	x (31, 49)
<i>H. sangarensis</i>	x (31)	
<i>Ixostrobus laxus</i>	x (20)	
<i>Juniperites gracilis</i>		x (54)
<i>Karkenia</i>	x (20)	
<i>Leptostrobus marginatus</i>	x (31)	x (31)
<i>Libocedrus arctica</i> (11)	x (20)	[3]
<i>L. (Andrivettia)</i>	x (24)	
<i>Mesocyparis beringiana</i>	x (31)	

<i>M. borealis</i>	x (31)
<i>Metasequoia cuneata</i>	x (31)
<i>M. disticha</i>	x (31)
<i>M. occidentalis</i>	x (31) x (31)
<i>Microconium beringiana</i>	x (18)
<i>Neozamites verchojanensis</i>	x (31)
<i>Nilssonia alaskana</i>	x (23, 24, 31) x (31, 32)
<i>N. comtula</i>	x (31)
<i>N. decursiva</i>	x (31, 49) x (31, 49)
<i>N. aff. grossinervis</i>	x (31)
<i>N. magnifolia</i>	x (31, 49) x (31, 49)
<i>N. menneri</i>	x (49) x (31, 49)
<i>N. pilifera</i>	x (31)
<i>N. schaumburgensis</i>	x (31)
<i>N. serotina</i>	x (31) x (31, 32, 33, 49, 54)
<i>N. cf. yukonensis</i>	x (31) x (31)
<i>Nilssoniocladus alaskensis</i>	x (50)
<i>N.chukotensis</i>	x (31, 50)
<i>Nilssoniopteris pryadae</i>	x (31)
<i>Otozamites jarmolenkoi</i>	x (24)
<i>Pagiophyllum triangulare</i>	x (31, 49) x (31, 49)
<i>Parataxodium cf. jacutensis</i>	x (31)

<i>P. neosibiricum</i>	x (20)	
<i>P. sugoijensis</i>	x (31)	
<i>P. wigginsii</i>	x (31, 49)	x (31, 49)
<i>Phoenicopsis angustifolia</i>	x (31, 20, 49)	x (31, 49)
<i>P. speciosa</i>	x (20)	x (31)
<i>Pinus</i> (6)	x (24)	x (55)
<i>P.</i> (?) sp.		x (32)
<i>Pityolepis tollii</i>	x (20)	
<i>Pityocladus</i>	x (31)	
<i>Pityophyllum arcticum</i>	x (20)	
<i>P. nordenskioldi</i>	x (20, 31, 49)	x (31, 49)
<i>P. pseudolaricis</i>	x (31)	
<i>P. staratschini</i>	x (20, 31, 49)	x (31, 49)
<i>Pityospermum brachypterum</i>	x (31)	[3]
<i>P. nansenii</i>	x (31)	
<i>P. pachypterum</i>	x (31)	
<i>P. aff. piniformis</i>	x (31)	
<i>P. semiovale</i>	x (31)	
<i>P. umbonatum</i>	x (31)	
<i>P. yukonense</i>	x (31)	
<i>Pitystrobus piceoides</i>	x (31, 49)	x (31, 49)
<i>P.</i> sp.		x (54)
<i>Podocarpus</i> (12)	x (24)	

<i>Podozamites corbinensis</i>	x (31)	
<i>P. eichwaldii</i>	x (20, 49)	x (31, 49)
<i>P. lanceolatus</i>	x (31)	x (31)
<i>P. latifolius</i>	x (20)	
<i>P. minor</i>	x (31)	
<i>P. aff. tuvensis</i>	x (31)	
<i>Protophyllocladus obesus</i>		x (32)
<i>P. aff. sachlinensis</i>	x (31)	
<i>Pseudocycas</i>	x (24)	x (31, 49)
<i>Pseudolarix arctica</i>	x (31)	[3]
<i>P. schmalhausenii</i>	x (20)	
<i>Pterophyllum validum</i>	x (31)	x (32)
<i>Sachalinella kolymensis</i>	x (31)	
<i>Sagenopteris variabilis</i> (Caytoniales)	x (31)	x (32)
<i>S. sp.</i>	x (24, 31)	x (31)
<i>Schizolepis cretaceous</i>	x (31)	
<i>Sciadopitys</i> sp.	x (31)	[3]
<i>Sequoia ambigua</i>		x (31)
<i>S. concinna</i>	x (31)	x (31)
<i>S. fastigata</i>	x (31)	x (31)
<i>S. cf. minuta</i>	x (31)	x (31)
<i>S. obovata</i>	x (31)	x (31)
<i>S. rigida</i>	x (31)	x (31)

<i>S. reichenbachii</i>	x (24, 31)	
<i>S. tenuifolia</i>	x (20, 31)	x (31)
<i>Sequoites artus</i>		x (54)
<i>S. dakotensis</i>		x (54)
<i>S. langsdorffii</i>		x (54)
<i>Sphenobaiera biloba</i>	x (31, 49)	x (31, 49)
<i>S. czeckanowskiana</i>	x (31, 49)	x (31, 49)
<i>S. flabellata</i>	x (20, 31)	x (31)
<i>S. longifolia</i>	x (2, 31, 49)	x (31, 49)
<i>S. pulchella</i>	x (31)	
<i>S. vera</i>	x (31)	
<i>Stenorachis</i> (Gymnospermae, incertae sedis), <i>S. striolatus</i>		x (31)
<i>S. sp.</i>	x (20)	
<i>Taeniopteris lundgreni</i>	x (31)	x (31)
<i>T. platyrachis</i>	x (31)	x (31)
<i>Taiwania microphylla</i>	x (20)	
<i>T. sp.</i>	x (24)	x (31)
<i>Taxites cephalotaxifolius</i>	x (31)	[3]
<i>T. olrikii</i>	x (31)	
<i>T. (Florinia?) sp.</i>	x (31)	
<i>Taxodium dubium</i>	x (31)	
<i>T. sp.</i>		x (32)

<i>Taxus</i>	x (24)	[3]
<i>Thuites intermedius</i>		x (33)
<i>T. interruptus</i>		x (54)
<i>T. sp.</i>	x (24)	
<i>Thuja cretacea</i>	x (31, 20)	x (31)
<i>T. interrupta</i>		x (33)
<i>Tollia cunninghamoides</i>	x (20)	
<i>Torellia anadyrensis</i>	x (31)	
<i>Torreya gramillima</i>		x (31)
<i>Torreyites tyrellii</i>		x (54)
<i>Williamsonia philatovii</i> (13)	x (31)	x (see 13)

Angiosperms

<i>'Acer' arcticum</i>	x (24)	x (33)
<i>A. collieri</i>		x (32)
<i>"A." quercifolium</i>	x (20)	
<i>Amaamia tschucotica</i>	x (31)	
<i>Anacardites neuburgae</i>	x (24)	[3]
<i>Anadyricarpa altingiosimila</i>	x (31)	
<i>Anona robusta</i>		x (54)
<i>Antholithes marsilioides</i>		x (54)
<i>Araliaecarpum kolymensis</i>	x (31)	
<i>Araliaephylloides arenaria</i>	x (31)	

<i>A. dentatum</i>	x (31)	
<i>A. devjatilova</i>	x (31)	x (31)
<i>A. cf. jarmoljukii</i>		x (31)
<i>A. leonvii</i>	x (31)	
<i>A. medium</i>	x (31)	x (31)
<i>A. pekulneense</i>	x (31)	
<i>A. pentalobatum</i>	x (31)	
<i>A. polevoii</i>	x (15, 31)	
<i>A. quinquelobum</i>	x (31)	x (31)
<i>A. speciosum</i>	x (31)	
<i>A. cf. subitum</i>	x (31)	x (31)
<i>Araliopsis</i>	x (24)	
<i>Archeampelos mullii</i>		x (31)
<i>Arctoterum rubifolium</i>	x (31)	
<i>Aristolochia paigei</i>	[52]	x (32)
<i>Aristolochites kamchaticus</i>	x (31)	[3]
<i>Arthollia inordinata</i>	x (15, 31)	(31)
<i>A. insignis</i>	x (31)	
<i>A. inundata</i>	x (31)	
<i>A. pacifica</i>	x (31)	x (31)
<i>A. rarytkensis</i>	x (31)	
<i>Arundo</i>	x (24)	[3]
<i>Barykonia kamchatica</i>	x (31)	

<i>B. tchucotica</i>	x (31)
<i>Caricopsis compacta</i>	x (31)
<i>C. laxa</i>	x (31)
<i>Castalites acutidentatus</i>	x (31, 32)
<i>C. flabelliformis</i>	x (31)
<i>C. inordinatus</i>	x (31)
<i>Celastrinites insignis</i>	x (33)
<i>C. septentrionalis</i>	x (31)
<i>C. wardii</i>	x (31)
<i>C. zakharovii</i>	x (31)
<i>C. sp.</i>	x (31)
<i>Celastrophyllum acutidens</i>	x (31)
<i>C. aff. kolymensis</i>	x (31)
<i>C. latifolium</i>	x (31)
<i>C. oppositifolia</i>	x (31)
<i>C. ovale</i>	x (24, 31)
<i>C. serrulatus</i>	x (31)
<i>C. sp.</i>	x (15, 24) x (31)
<i>Celastrus herendeensis</i>	x (31)
<i>C. pseudocurvinervis</i>	x (31)
<i>C. septentrionalis</i>	x (18)
<i>C. sp.</i>	x (15)
<i>Celtis</i>	x (24) [3, w]

<i>Cercidiphyllum potomacensis</i>	x (31)	[3, w]
<i>Cinnamomoides ievlevii</i>	x (31)	[<i>Cinnamomum</i>
		w]
<i>Cissites(?) asymmetricus</i>	x (31)	
<i>C. beljaevii</i>	x (31)	x (31, 32)
<i>C. comparabilis</i>	x (24)	
<i>C. cordatus</i>		x (31)
<i>C. hermanii</i>	x (31)	
<i>C. inscissus</i>	x (31)	
<i>C. kautajamensis</i>	x (31)	
<i>C. microphylla</i>	x (24)	
<i>C. orbiculatus</i>	x (31)	x (31)
<i>C. pekulneensis</i>	x (31)	
<i>C. volobuevae</i>	x (18)	
<i>Cissus pseudomarginata</i>	x (31)	[3, w]
<i>Cocculophyllum cf. furcinerve</i>	x (31)	
“ <i>Cocculus</i> ” cf. <i>extinctus</i>	x (31)	
<i>C.</i> (?) sp.		x (31)
<i>Corylites beringianus</i>	x (31)	x (31)
<i>Corylus ageevii</i>	x (31)	
<i>C. beringiana</i>	x (18)	
<i>C. denverensis</i>		x (33)
<i>Crataegites cf. borealis</i>	x (24, 31)	[3, w]

“ <i>Credneria</i> ” cf. <i>grewiopsoides</i>	x (31, 32)
<i>Credneria</i> (= <i>Artholia</i>)	x (20, 24, 31)
<i>C.</i> sp.	x (31)
<i>Dalbergites elegans</i>	x (31)
<i>D. cf. simplex</i>	x (31)
<i>Dalembia farajevii</i>	x (31)
<i>D. krassilovii</i>	x (31)
<i>D. pergamentii</i>	x (15, 31) x (31)
<i>D. vachrameevii</i>	x (23, 31) x (31)
<i>Daphnophyllum dakotense</i>	x (32)
<i>D.</i> sp.	x (24)
“ <i>Diospyros</i> ” aff. <i>steenstrupi</i>	x (31) x (31, 32)
<i>D.</i> sp.	x (24)
<i>Disanthophyllum longipetiolatum</i>	x (31)
<i>Dombeyopsis nebrascensis</i>	x (54)
<i>Dyrana orientalis</i>	x (31)
<i>Ettinghausenia affinis</i>	x (31)
<i>E. embicola</i>	x (31)
<i>E. cf. latilobata</i>	x (31)
<i>E. louravetlanica</i>	x (31) x (31)
<i>E. cf. marginata</i>	x (31)
<i>E. newberryana</i>	x (31)
<i>E. primaeva</i>	x (31) x (31)

<i>E. prisca</i>	x (31)	
<i>E. rarivervis</i>	x (31)	
<i>E. raynoldsii</i>	x (31)	x (31)
<i>Fraxinus leii</i>		x (54)
<i>Grebenia anadyrensis</i>	x (31)	x (31)
<i>Grewiopsis akhmetjevii</i>	x (31)	
<i>G. nemorosus</i>	x (31)	
<i>G. yukonensis</i>		x (32)
<i>Haemanthopodium cordatum</i>	x (16, 18, 31)	
<i>H. kamtschaticum</i>		x (31)
<i>Hedera curvata</i>		x (32)
<i>H. gracile</i>	x (20)	
“ <i>Hedrea</i> ” sp.		x (31)
<i>Hollickia quercifolia</i>	x (31)	
<i>Ievlevia dorofeevii</i>	x (31)	
<i>Ilex vernicosa</i> (14)	x (31)	[3, w]
<i>Jenkinsella arctica</i>		x (54)
<i>Juglans lecontiana</i>		x (33)
<i>J. sp.</i>	x (24)	
<i>Kakanautia repanda</i>	x (31)	
<i>Kenella filatovii</i>	x (31)	x (31)
<i>K. harrisiana</i>	x (31)	
<i>Laurophyllo laraminum</i>		x (33)

<i>Laurophylloides (Laurus plutonia)</i>	x (24)
<i>Leguminosites stagnum</i>	x (54)
<i>Limnobiophyllum scutatum</i>	x (16)
<i>Lindera jarmolenkoi (Sassafras polevoii)</i>	x (24)
<i>L. minima</i>	x (31) [3]
<i>Liriophyllum aeternum</i>	x (31) x (31)
<i>Macclintockia beringiana</i>	x (31)
<i>M. crenata</i>	x (31)
<i>M. decurens</i>	x (20)
<i>M. cf. dentata</i>	x (31)
<i>M. electilis</i>	x (32)
<i>M. neosibirica</i>	x (20)
<i>M. ochotica</i>	x (23, 31)
<i>M. sp.</i>	x (15, 24)
<i>Magnolia amplifolia</i>	x (31)
“ <i>Magnolia</i> ” cf. <i>lakesii</i>	x (31)
<i>M. sp.</i>	x (24)
<i>Magnoliaeophyllum alternans</i>	x (31) x (31)
<i>M. magnificum</i>	x (31)
<i>Menispermites belli</i>	x (33)
<i>M. efimovae</i>	x (31)
<i>M. ginterensis</i>	x (31)
<i>M. grebenkaens</i>	x (31)

<i>M. marcovoensis</i>	x (31)	
<i>M. minutus</i>	x (31)	
<i>M. aff. reniformis</i>	x (31)	x (31)
<i>M. septentrionalis</i>	x (15, 31)	x (31)
<i>M. sibiricus</i>	x (31)	x (31)
<i>M. vassetkii</i>	x (31)	
<i>Mirella borealis</i>	x (31)	
<i>Myrtophyllum acuminata</i>	x (31)	
<i>M. penzhinense</i>	x (31)	
<i>Myrica(?) trifoliata</i>		x (32)
<i>M. sp.</i>	x (24)	
<i>Nelumbites aff. minimus</i>	x (31)	
<i>N. protoluteus</i>		x (33)
<i>N. sp.</i>	x (24)	x (31)
<i>Nelumbo</i>	x (24)	
<i>Nordenskioldia</i>	x (24)	[3]
<i>Nuphar</i>	x (31)	
<i>Nymphaeites angulatus</i>		x (33, 54)
<i>N. striatus</i>		x (54)
<i>N. sp.</i>	x (31)	
<i>Nyssa tschucotica</i>	x (18)	[3]
<i>Nyssidium arcticum</i>	x (31)	x (31)
<i>Palaeotrapa aculeata</i>	x (31)	

<i>P. bicornata</i>	x (31)	
<i>P. triangulata</i>	x (31)	
<i>Paliurus aff. visibilis</i>	x (31)	x (32)
<i>Paraprotophyllum cordatum</i>	x (31)	
<i>P. ignatianum</i>	x (24, 31)	x (31)
<i>P. pseudopeltatum</i>	x (31)	
<i>Peculnea lancea</i>	x (31)	
<i>P. pinnatiloba</i>	x (31)	
<i>Penzhinia vachrameevii</i>	x (31)	
<i>Phragmites alaskana</i>	x (31)	x (31)
<i>Pistia corrugata</i>		x (33)
<i>Pistia(?) marginata</i>	x (24)	
<i>Platanophyllum</i> sp.		x (54)
<i>Platanus basilobata</i>		x (33)
<i>Platanus cuneifolia</i>	x (24)	
<i>P. cuneiformis</i>	x (24)	
<i>P. embicolia</i>	x (23, 24)	
<i>P. latior</i>		x (32)
<i>P. louravetlanica</i>	x (49)	x (49)
<i>P. reynoldsi</i>	x (18, 31)	x (54)
<i>P. sp.</i>	x (15, 23, 24, 31)	x (31)
<i>Platimelis platanoides</i>	x (31)	
<i>P. umcuvensis</i>	x (31)	

<i>Populites</i> cf. <i>pseudolancastriensis</i>	[52]	x (31)
<i>Potomageton</i>	x (24)	
<i>Protophyllum boreale</i>		x (31)
“ <i>Protophyllum</i> ” <i>terechova</i>	x (31)	
<i>Pseudoprotophyllum boreale</i>	x (20, 31)	x (31)
<i>P. dentata</i>	x (31)	
<i>P. giganteum</i>	x (20)	
<i>P. parvaefolium</i>	x (20)	
<i>P. sp.(2)</i>	x (15, 20, 24)	
<i>Quercus groenlandica</i>	x (31)	
<i>Q. pseudomarionii</i>		x (32)
<i>Q. tchucotica</i>	x (15, 23)	
“ <i>Quercus</i> ” <i>viburnifolia</i>	x (31)	
<i>Quereuxia</i>	x (15, 23, 24)	
<i>Q. angulata</i>	x (16, 24, 31)	x (31)
<i>Q. flabellata</i>	x (31)	
<i>Q. rotundifolia</i>	x (31)	
<i>Rarytkinia amaamensis</i>	x (31)	
<i>R. aff. terechovae</i>	x (18, 31)	x (31)
<i>Rhamnites cornifolia</i>		x (32)
<i>R. cf. eminens</i>	x (31)	
<i>R. septentrionalis</i>	x (31)	
<i>Renea actinidifolia</i>	x (31)	

“ <i>Rubus</i> ”	x (31)	x (31)
<i>Samaropsis cirrifera</i>	x (31)	
<i>Sapindopsis</i>	x (31)	
<i>Sapotacites</i> cf. <i>alaskensis</i>	x (31)	[3]
<i>Scheffleraephyllum venustum</i>	x (31, 49)	x (31, 49)
<i>Smilax</i> aff. <i>grandifolia</i>	x (31)	
<i>S.</i> sp.		x (31)
<i>Sorbus alaskana</i>		x (32)
<i>Sorbus(?)</i> sp.	x (18)	[3]
<i>Sparganiophyllum multinervosum</i>	x (16)	
<i>S.</i> sp.	x (31)	x (31)
<i>Spirodela scutata</i>		x (54)
<i>Stenorachis lepida</i>	x (31)	
<i>Sterculia basiauriculata</i>		x (32)
<i>S.</i> sp.	x (24)	
<i>Sugio opposita</i>	x (31)	
<i>Terechovia anadyrensis</i>	x (31)	
<i>T. intermedia</i>	x (31)	
<i>Ternstroemites harwoodensis</i>	x (15, 31)	[3]
<i>Tiliaeephyllum (Tilia) brooksense</i>		x (31)
<i>T. tsagaianicum</i>	x (24)	
<i>Trochodendrocarpus</i>	x (24)	
<i>Trochodendroides arctica</i>	x (20, 31)	x (31, 54)

<i>T. bidentata</i>	x (18, 31)
<i>T. bifida</i>	x (31)
<i>T. emimensis</i>	x (31)
<i>T. grossidentata</i>	x (31)
<i>T. lamutensis</i>	x (18, 31) x (31)
<i>T. longifolia</i>	x (31)
<i>T. notabilis</i>	x (23, 31) x (31)
<i>T. pekulnejensis</i>	x (31)
<i>T. richardsonii</i>	x (20, 31)
<i>T. sachalinensis</i>	x (31)
<i>T. smilacifolia</i>	x (20)
<i>T. speciosa</i>	x (31)
<i>T. vassilenkoi</i>	x (31) x (31)
<i>Typha</i>	x (54)
<i>Ulmus oblongifolia</i>	x (31)
<i>U. sp.</i>	x (31)
<i>Viburniphyllyum alnoides</i>	x (31)
<i>V. basicordatum</i>	x (31)
<i>V. lebedevii</i>	x (31)
<i>V. cf. newberrianum</i>	x (31)
<i>V. whymperi</i>	x (15, 31) x (31)
<i>Viburnum acutifolium</i>	x (31)
<i>V. aff. asperum</i>	x (31) x (33)

<i>V. oblongum</i>	x (31)	
<i>V. pseudoantiquum</i>	x (31)	
<i>V. rarytkense</i>	x (31)	x (31)
<i>V. simile</i>		x (33, 54)
<i>V. stenophyllum</i>	x (31)	
<i>Vitis dakotana</i>		x (33)
<i>Vitis olrikii</i>	x (31)	
“ <i>Vitis</i> ” <i>penzhinica</i>	x (31)	
“ <i>V.</i> ” <i>populoides</i>	x (31)	x (32)
“ <i>V.</i> ” <i>stantoni</i>	x (31)	x (33, 54)
<i>Vitis</i> (24, p. 212)	x (24)	
<i>Zizyphoides colombi</i>		x (33)
<i>Z. heterophylla</i>	x (24)	
<i>Z. mackayi</i>		x (33)
<i>Z. smilacifolia</i>	x (20)	
<i>Z. varietas</i>	x (20)	
<i>Zizyphus anadyrensis</i>	x (31)	
<i>Z. electilis</i>	x (31)	
“ <i>Z</i> ”. <i>kuijensis</i>	x (31)	x (31)
“ <i>Z</i> .” <i>smilacifolia</i>	x (31)	x (31)

Paleocene

Ferns and allied groups

<i>Amurocyparis</i>	x (29)	
<i>Asplenium? penhallowi</i>	x (54)	
<i>Asplenium</i> (a)	x (29)	[3]
<i>Azolla schopfii</i>	x (53)	
<i>A.vellus</i>	x (53)	
<i>Cladophlebis cf. arctica</i>	x (24)	[3]
<i>C. groenlandica</i>	x (54)	
<i>C. oerstedtii</i>	x (24)	
<i>Dennstastra sorimarginata</i>	x (53)	
<i>Dennstaedtia americana</i>	x (35, 53)	
<i>D. blomstrandia</i>	x (54)	
<i>D. sp.</i>	x (29)	
<i>Dryopterites georgii?</i>	x (54)	
<i>Equisetum alexoensis</i>	x (54)	
<i>E. arctium</i>	x (54)	
<i>E. boreale</i>	x (54)	
<i>E. fluviatoides</i>	x (54)	
<i>E. sp.</i>	x (22, 29)	[3]
cf. <i>Isöetites horridus</i>	x (53)	
<i>Onoclea hebridica</i>	x (54)	
<i>Onoclea hespera</i>	x (35, 53)	
<i>O. sp.</i>	x (29)	
<i>Osmunda macrophylla</i>	x (35, 53, 54)	

<i>O. sachalinensis</i>	x (24)	
<i>Salvinia</i>	x (29)	[3]
<i>Woodwardia gravida</i>		x (53)
<i>W. sp.</i>	x (24)	[3]

Gymnosperms

<i>Androvettia catenulata</i>	x (54)	
<i>Chamaecyparis</i>	x (29)	[3]
<i>Cryptomerites lambii</i>		x (54)
<i>Cupressinocladius</i> (a)	x (29)	[3]
<i>Elatocladus campbellii</i>		x (53)
<i>E. nordenskioldi</i>		x (54)
<i>E. olriki</i>		x (54)
<i>E. tinajorum</i>		x (54)
<i>E. ungeri</i>		x (54)
<i>Fokienia ravenscragensis</i>		x (53)
<i>Fokiensiopsis</i> (a)	x (29)	
<i>Ginkgo adianoides</i>	x (21, 24)	
<i>G. biloba</i>		x (35)
<i>G. cretaceae</i>	x (24)	
<i>G. spitsbergensis</i>		x (53)
<i>Ginkgoites adiantoides</i>		x (54)
<i>Glyptostrobus europaeus</i> (b)	x (24)	

<i>G. nordenskioldi</i>	x (35, 53)
<i>Mesocyparis borealis</i>	x (53)
<i>Metasequoia foxii</i>	x (36)
<i>M. occidentalis</i>	x (21, 22) x (53, w)
<i>Quasisequoia couttsiae</i>	x (29)
<i>Sequoiites artus</i>	x (54)
<i>S. dakotensis</i>	x (54)
<i>S. langsdorffii</i>	x (54)
<i>Taxites olrikii</i> (b)	x (29) [3]
<i>Taxodium dubium</i>	x (24) x (w)
<i>Thuites interruptus</i>	x (54)

Angiosperms

<i>Acer arcticum</i> (a)	x (21)	[3], (54)
<i>Aesculus magnificum</i>	x (21)	x (40)
<i>Alnus beringiana</i>	x (21)	
<i>A. serrata</i>		x (54)
<i>A. sp.</i>	x (24, 29)	x (35)
<i>Amelanchites similis</i>		x (53)
<i>Anoectomeria brongniartii</i>	x (29)	
<i>Apollonias</i>	x (29)	
<i>Aralia alexoensis</i>		x (54)
<i>A. venulosa</i>	x (29)	[3]

<i>Archeampelos acerifolia</i>	x (53)	
<i>A. mullii</i>	x (51)	
<i>Artocarpus</i>	x (29)	[3]
<i>Betula</i> (b)	x (29)	[3]
<i>Bumelia oblongifolia</i>	x (29)	[3]
<i>Callistemophyllum melaleucaeformae</i>	x (29)	[3]
<i>Carex</i> (b)	x (29)	[3]
<i>Carpinus</i> (b)	x (29)	[3]
<i>Celastrinites insignis</i>		x (54)
<i>Cornophyllum newberryi</i>		x (53)
<i>C. tenuinervia</i>	x (21)	
<i>Cornus denverensis</i>		x (54)
<i>C. krassilovi</i>	x (14)	
<i>Corylites</i>	x (29)	[3]
<i>C. fosteri</i>	x (35)	x (54)
<i>Corylus</i> (a)	x (29)	[3]
<i>Craigia</i> (b)	x (29)	
<i>Cryptomerites</i>	x (29)	
<i>Cyperacites</i>	x (29)	[3]
<i>Daphnogene lanceolata</i>	x (29)	[3]
<i>Davidia</i> (b)	x (29)	[3]
<i>Dewalquea gelindensis</i>	x (29)	
<i>Dombeyopsis nebrascensis</i>		x (54)

<i>Dryandra karakulensis</i>	x (29)	[3]
<i>Dryophyllum</i> sp.		x (54)
<i>Fagopsis</i> (b)	x (29)	[3]
<i>Ficus murczisonii</i> (b)	x (29)	[3]
<i>Hydrangea</i> (b)	x (29)	[3]
<i>Jenkinsella arctica</i>		x (54)
<i>Juglans lecontiana</i>		x (54)
<i>J. nigella</i>		x (54)
<i>J. thermalis</i>		x (54)
<i>Juglans</i> (a)	x (29)	[3]
<i>Laurophylum laraminum</i>		x (54)
<i>L.</i> sp.	x (29)	[3]
<i>Laurus</i> (b)	x (29)	[3]
<i>Leucothoë protogaea</i> (b)	x (29)	[3, Q]
<i>Limnobiophyllum scutatum</i>	x (16)	x (17)
<i>Lindera</i> (a)	x (29)	[3]
<i>Litsea</i>	x (29)	[3]
<i>Macclintockia</i> (a)	x (29)	[3]
<i>Magnolia embica</i> (a)	x (29)	[3]
<i>Menispermites belli</i>		x (54)
<i>Mimosites</i>	x (29)	[3]
<i>Myrica banksiaeefolia</i> (a)	x (29)	[3]
<i>M. beringiana</i>	x (22)	

<i>Myrciophyllum americanum</i>	x (53)	
cf. <i>Myristica catahouensis</i>	x (53)	
<i>Nelumbo montanum</i>	x (53)	
<i>Nelumbites protoluteus</i>	x (54)	
<i>Nelumbo</i> (b)	x (29)	[3]
<i>Nymphaea polyrhiza</i>	x (29)	[3]
<i>Nymphaeites angulatus</i>		x (54)
<i>N. striatus</i>		x (54)
<i>Nyssa bureica</i> (b)	x (29)	[3]
<i>Nyssidium</i> (a)	x (29)	[3, as <i>Cercidiphyllum</i>]
<i>Ocotea</i> (a)	x (29)	[3]
<i>Oxycarpia bifaria</i>	x (29)	
<i>Palaeocarpinus</i>	x (29)	
<i>Paranymphaea crassifolia</i>		x (53, 54)
<i>Peltandra</i>	x (29)	[3]
<i>Persea enormis</i> (b)	x (29)	[3]
<i>P. paleomorpha</i>	x (29)	
<i>Phoebe</i> (b)	x (29)	[3]
<i>Phragmites</i> (a)	x (29)	[3]
<i>Pistia corrugata</i>		x (53, 54)
<i>Platanites canadensis</i>		x (53)
<i>Platanus basicordata</i> (a)	x (21)	[3]

<i>P. basilobata</i>	x (54)	
<i>P. raynoldsii</i>	x (54)	
<i>Poacites</i>	x (29)	[3]
<i>Populus carneosa</i>	x (54)	
<i>P. daturaefolia</i>	x (54)	
<i>P. penhallowi</i>	x (54)	
<i>P. raynoldsii</i>	x (29)	
<i>P. snatolensis</i> (b)	x (21)	[3]
<i>P. sp.</i>	x (22, 24)	
<i>Porosia</i>	x (29)	
<i>Protoacerophyllum perfoliatum</i>	x (29)	
<i>Protophyllum</i> (a)	x (24)	[3]
<i>Prunites maclearnii</i>	x (53)	
<i>Pterocarya</i> (b)	x (29)	[3]
<i>Pterospermites dawsoni</i>	x (54)	
<i>P. haguei?</i>	x (54)	
<i>P. whitei</i>	x (54)	
<i>Quercus praegroenlandica</i>	x (54)	
<i>Rhamnites cleburnii</i>	x (53)	
<i>R. marginatus</i>	x (54)	
<i>R. ovatus</i>	x (54)	
<i>Rorippides aquatica</i>	x (53)	
“ <i>Sapindus</i> ” <i>affinis</i>	x (35)	[3]

<i>cf. Sapindus grandifoliolus</i>		x (53)
<i>S. sp.</i>	x (29)	
<i>Sideroxylon</i> (b)	x (29)	[3]
<i>Sorbus</i> (a)	x (29)	[3]
<i>Spirodela scutata</i>		x (54)
<i>Tetracentron</i> (b)	x (29)	
<i>Tiliaeophyllum brookense</i>		x (51)
<i>T. sp.</i>	x (29)	
<i>Trapago angulata</i>		x (53)
<i>Trochodendrocarpus</i>	x (29)	
<i>Trochodendroides arctica</i>	x (21, 22, 24,	x (35, 54)
	29)	
<i>T. flabellata</i>		x (53)
<i>T. serrulata</i>		x (35)
<i>T. speciosa</i>		x (53)
<i>Ulmus furcinervis</i>	x (29)	[3]
<i>U. cf. longifolia</i>	x (24)	
<i>U. pseudobraunii</i>	x (21, 24)	
<i>Ushia alnifolia</i>	x (29)	
<i>U. janishevskii</i>	x (29)	
<i>U. kamyschinensis</i>	x (29)	
<i>Viburnum antiquum</i> (a)	x (21)	[3], x (54)
<i>V. asperum</i>		x (54)

<i>V. lakesii</i>	x (54)
<i>V. ravenscragensis</i>	x (53)
<i>V. simile</i>	x (54)
<i>V. trinervum</i>	x (54)
<i>V. volgensie</i>	x (29)
<i>Vitis stantonii</i>	x (54)
<i>V. sp.</i>	x (53)
<i>Zingiberopsis</i>	x (29)
<i>Zizyphoides colombi</i>	x (54)
<i>Z. mackayi</i>	x (54)

Eocene

Ferns and allied groups

<i>Acrostichum</i>	x (29)	[3]
<i>Anemia gunni</i>		x (37)
<i>Anemia</i>	x (29)	
<i>Azolla vera</i>	x (29)	[3]
<i>Blechnum akartscheschmens</i>	x (29)	[3]
<i>Cyclosorus</i>	x (29)	
<i>Dennstaedtia americana</i>	x (1, 21, 22)	x (3, w)
	29)	
<i>Dryopteris alaskana</i>		x (37)
<i>D. grosseserrata</i>	x (1, 21)	

<i>Equisetum arcticum</i>	x (1, 21, 29)	x (w, 3)
<i>Hydropteris indutus</i> (3)	x (29)	
<i>Lastrea badchyzica</i>	x (29)	[3]
<i>Lygodium asiaaticum</i>	x (29)	
<i>L. coloradense</i>	x (29)	
<i>L. larsoni</i>		x (37)
<i>Onoclea hesperia</i> (a)	x (29)	[3]
<i>Osmunda macrophylla</i>	x (1, 21, 22,	x (BC, 2)
	29)	
<i>O. sachalinensis</i>	x (29)	
<i>Osmundastrum heislianum</i>	x (1, 21, 22)	
<i>Polypodites nepphrotiopsis</i>	x (29)	[3]
<i>Regnellidium</i>	x(29)	
<i>Salvinia mildeana</i> (b)	x (29)	[3]
<i>S. natanella</i>	x (29)	
<i>Woodwardia</i> (b)	x (29)	[3]

Gymnosperms

<i>Abies tigilensis</i> (b)	x (1, 21)	[3]
<i>Araucaria dubia</i>	x (1, 21)	
<i>Athrotaxis</i>	x (29)	[3]
<i>Cryptomeria kamtschatica</i> (a)	x (1, 21)	
<i>Cunninghamia sinelnikovae</i> (a)	x (1, 21)	[3]

<i>Doliostrobus</i>	x (29)	
<i>Ginkgo adiantoides</i> (a)	x (1, 21, 29)	[3]
<i>G. kamtschatica</i>	x (22, 29)	
<i>Glyptostrobus europaeus</i>	x (1, 29)	x (5, w)
<i>G. groenlandicus</i>	x (21)	
<i>G. nordenskioldii</i>	x (22)	
<i>G. sp.</i>		x (37)
<i>Larix</i> (b)	x (29)	[3]
<i>Metasequoia glyptostroboides</i>		x (37)
<i>M. occidentalis</i>	x (1, 21, 22,	x (6, w)
	29)	
<i>Picea</i> sp. (b)	x (1, 29)	[3]
<i>Pinus kovatschensis</i>	x (1, 21)	[3]
“ <i>Podocarpus</i> ” <i>eocenica</i> (b)	x (29)	[3]
<i>Pseudolarix</i> (b)	x (29)	[3]
<i>Quasisequoia couttsiae</i>	x (29)	
<i>Sciadopitys sveshnikovae</i> (b)	x (1, 21)	[3]
<i>Sequoia affinis</i>	x (1, 21)	x (7, w)
<i>Taxodium dubium</i>	x (1, 21, 29)	x (5, w)
<i>T. tinajorum</i>	x (29)	
<i>Thuites napanensis</i> (a)	x (1, 21)	[3]
<i>Torreya kamschatcica</i>	x (1)	[3]
<i>T. seminovii</i>	x (21)	

Angiosperms

<i>Acer arcticum</i>	x (1, 21, 22)	x (11)
<i>A. ezoanum</i>	x (29)	
<i>A. glabroides</i>		x (37)
<i>A. kinkilense</i>	x (1, 21)	
<i>A. cf. kushiroanum</i>	x (29)	
<i>A. cf. palaeoplatanoides</i>	x (29)	
<i>Actinidia ovata</i> (b)	x (21)	
<i>A. semenovii</i>	x (1)	
<i>Aesculus hickeyi</i>		x (40)
<i>A. magnifica</i>	x (22, 29)	
<i>Ailanthus</i> (b)	x (29)	[3]
<i>Alangium bergensis</i>		x (37)
<i>A. lenaense</i>	x (1, 21)	
<i>Alnus beringiana</i>	x (22)	
<i>A. cuprovallis</i>		x (37)
<i>A. ezoensis</i>	x (29)	
<i>A. hokkadoensis</i>	x (29)	
<i>A. ishikariensis</i>	x (29)	
<i>A. kushiroensis</i>	x (29)	
<i>A. onorica</i>	x (29)	
<i>A. protophylloides</i>	x (1, 21, 29)	

<i>A. raridentata</i>	x (1)	
<i>Ampelopsis cercidifolia</i> (b)	x (1)	[3]
<i>A. cordifolia</i>	x (21)	
<i>A. schischkinii</i>	x (29)	
<i>Amygdalus turcmensis</i> (b)	x (29)	[3]
<i>Anacardites badchyzicus</i>	x (29)	[3]
<i>Andromeda</i>	x (29)	[3]
<i>Apocynophyllum helveticum</i>	x (25)	[3]
<i>Archeampelos acerifolia</i>	x (29)	
(= <i>Acer arcticum</i> fide 29)		
<i>Arunidites roginei</i>	x (29)	[3]
<i>Asclepiadites</i>	x (21)	[3]
<i>Astronium ninae</i> (b)	x (29)	[3]
<i>Banksia myricifolia</i> (b)	x (29)	[3]
<i>Betula acutidentata</i>	x (1, 21)	
<i>B.</i> sp.	x (37)	
<i>Broussonetia kamtschatica</i> (b)	x (1, 21)	
<i>Bytneria iizimae</i> (b)	x (29)	
<i>Camellia</i> (b)	x (29)	
<i>Carpinus japonica</i> (4)	x (29)	x (w)
<i>C. kushiroensis</i>	x (29)	
<i>Carya atlantica</i>	x (21)	
<i>C. casmanensis</i>	x (37)	

<i>C. grandidentata</i>	x (21)	
<i>C. typical</i>	x (29)	
<i>Cassia</i> (b)	x (29)	[3]
<i>Castanea kamtschatica</i> (b)	x (1, 21, 29)	[3]
<i>C. longifolia</i>	x (29)	
<i>C. sachalinensis</i>	x (29)	
<i>Castanopsis kazachstanensis</i> (b)	x (29)	[3]
<i>C. pavlodarensis</i>	x (29)	
<i>Ceanothus rectinervus</i> (b)	x (1, 21)	[3]
<i>Cedrela</i> (b)	x (29)	[3]
<i>Celtis</i> (b)	x (29)	[3]
<i>Cercidiphyllum crenatum</i>		x (37)
<i>C. grandifolium</i>	x (1)	
<i>C. intermedium</i>	x (1, 21, 29)	
<i>C. kamtschaticum</i>	x (1, 21)	
<i>Cinnamophyllum kushtakensis</i>		x (37)
<i>C.</i> sp.	x (29)	
<i>Cissites ochotensis</i> (a)	x (29)	[3]
<i>Cissus beringiana</i> (b)	x (21)	[3]
<i>Coccoloba</i> (b)	x (29)	[3]
<i>Cocculus ezoanum</i>	x (29)	
<i>C.</i> sp.		x (37)
<i>Comptonia</i> (b)	x (29)	[3]

<i>Cordia ochotensis</i> (b)	x (21)	[3]
<i>Cornophyllum</i> sp.	x (1, 21, 22; see 14)	
<i>Corylopsiphyllum carpinifolium</i>	x (21)	
<i>Corylopsis kamtschatica</i>	x (21, 22)	
<i>C.</i> sp.		x (37)
<i>Corylus kinkilensis</i> (4)	x (1, 21)	x (w)
<i>C. palaeomaximovicziana</i>	x (29)	
<i>Cotinus</i>	x (29)	[3]
<i>Cragia oregonensis</i> (b)	x (29)	
<i>Crataegus</i> (b)	x (29)	[3]
<i>Daphne</i> (b)	x (29)	[3]
<i>Daphnogene</i>	x (29)	[3]
<i>Davidoidea kamschatica</i>	x (22)	
<i>Delavaya</i>	x (29)	
<i>Diospyros dubiosa</i> (a)	x (1)	[3]
<i>Dombeya</i> (b)	x (29)	
<i>Dryophyllum curticellense</i>	x (29)	
<i>D. dewalque</i>	x (29)	
<i>D. kryshtofovichii</i>	x (29)	
<i>D. pugetensis</i>		x (37)
<i>Echitonium</i>	x (29)	[3]
<i>Eotrigonobalanus</i>	x (29)	

<i>Eucalyptus</i> (b)	x (29)	[3]
<i>Eugenia haeringiana</i>	x (29)	
<i>E. kirshiensis</i>	x (29)	
<i>E. sp.</i>		x (37)
<i>Euonymus denticulata</i>	x (29)	[3]
<i>Euphorbiophyllum erodofolium</i>	x (21)	[3]
<i>E. insolitum</i>	x (1)	
<i>Fagopsis nipponica</i>	x (29)	
<i>F. undulata</i>	x (1)	x (9, 10)
<i>Fagus napanensis</i>	x (1, 21, 29)	
<i>F. sp.</i>		x (37)
<i>Ficus spiculifolia</i> (b)	x (29)	[3]
<i>Firmiana alaskana</i>		x (37)
<i>F. kovatschensis</i>	x (1, 21)	
<i>Fothergilla</i>	x (29)	x (37)
<i>Fraxinus</i> (b)	x (29)	[3]
<i>Grewiopsis auriculaecordatus</i>	x (21)	[3]
<i>G. conjunctivus</i>	x (21)	
<i>Guoania</i>	x (29)	
<i>Hakea spathulata</i> (b)	x (29)	[3]
<i>Hamamelites kinkilensis</i>	x (1)	[3]
<i>H. palanensis</i>	x (21)	
<i>H. snatolensis</i>	x (22)	

<i>H. viburnifolius</i>	x (21)	
<i>Hamamelis kushiroensis</i> (b)	x (29)	[3]
“ <i>Hibiscus</i> ” <i>kalmakpaica</i>	x (29)	[3]
“ <i>H.</i> ” <i>neuburgae</i>	x (29)	
<i>Hydrangea alnifolia</i> (b)	x (21, 22)	[3]
<i>Idesia romanovii</i> (b)	x (29)	[3]
<i>Juglans beringiana</i>	x (21)	
<i>J. palaeocathayensis</i>	x (22)	
<i>J. pogozhevii</i>	x (1, 21, 29)	
<i>J. protocinerea</i>	x (1, 29)	
<i>J. sp.</i>		x (37)
<i>Kalopanax latifolia</i> (b)	x (21)	
<i>Kleinhovia bastiruncata</i> (b)	x (29)	
<i>Koelreuteria kamtschatica</i> (b)	x (1, 21)	[3]
<i>Laurus</i> (b)	x (29)	[3]
<i>Leucothoë protogaea</i> (b)	x (29)	[3]
<i>Limnobiophyllum</i> (a)	x (29)	
<i>Lindera vassilenkoi</i> (a)	x (29)	
<i>Liquidambar brevilibata</i>	x (1, 29)	
<i>L. europaea</i>	x (29)	
<i>L. kazakhstanica</i>	x (29)	
<i>L. miosinica</i>	x (29)	
<i>L. sp.</i>		x (37)

<i>Lithocarpus</i> (b)	x (29)	[3]
<i>Litsea</i> (b)	x (29)	[3]
<i>Litseaphyllum carbonensis</i>		x (37)
<i>Lomatia macrophylla</i> (b)	x (29)	[3]
<i>Macclintockia pugetensis</i>	[52]	x (37)
<i>Macginitiea grandifolia</i>	x (1, 21)	
<i>M. kanei</i>	x (29)	
<i>Magnolia basicordata</i> (a)	x (22)	
<i>M. kryshtofovichii</i>	x (29)	
<i>M. ochotica</i>	x (21)	
<i>M. praekobus</i>	x (1, 21, 29)	
<i>M. reticulata</i>		x (37)
<i>Majanthemophyllum</i>	x (29)	[3]
<i>Melastomites</i>	x (29)	
<i>Mallotus hokkaidoensis</i> (b)	x (29)	[3]
<i>Musophyllum complicatum</i>	x (1, 21)	x [9, 12, 13 (as <i>Mussopsis)</i>]
<i>Myrica acuminata</i> (b?)	x (29)	[3]
<i>M. beringiana</i>	x (1, 21, 29)	
<i>M. dilodendrofolia</i>	x (29)	
<i>M. cf. lignitum</i>	x (29)	
<i>M. longifolia</i>	x (25)	
<i>M. ovalifolia</i>	x (29)	

<i>M. vindobonensis</i>	x (29)	
<i>Myrsine</i> (b)	x (29)	[3]
<i>Nelumbo nipponica</i> (b)	x (29)	[3]
<i>N. protospeciosa</i>	x (21)	
<i>Neolitsea</i>	x (29)	[3]
<i>Nitophyllites</i>	x (29)	
<i>Nordenskioldia borealis</i>	x (22, 29)	[3]
<i>N.</i> sp.	x (38, 39)	x (38, 39)
<i>Nyssidium arcticum</i> (a)	x (22, 29)	[3]
<i>Nuphar</i> (b)	x (29)	
<i>Nypa burtinii</i>	x (29)	(w, tropical)
<i>Nyssa dubiosa</i> (b)	x (21)	
<i>N. integerrima</i>	x (1)	
<i>Ocotea</i> (a)	x (29)	
<i>Oxycarpia</i>	x (29)	
<i>Palibinia pinnatifida</i>	x (25)	
<i>Peltandra</i> (b)	x (29)	[3]
<i>Persea</i> (b)	x (29)	[3]
<i>Phellodendron kovatschense</i>	x (21)	
<i>Phragmites</i>	x (22)	[3]
<i>Phytocrene acutissima</i>		x (37)
<i>P.</i> sp.	x (29)	
<i>Pittosporum</i>	x (29)	

<i>Platkeria basiobliqua</i>	x (29)
<i>Platanus acutiloba</i>	x (29)
<i>P. comstocki</i>	x (37)
<i>P. iljinskajae</i>	x (22)
<i>P. insolida</i>	x (21)
<i>P. mabuti</i>	x (29)
<i>P. newberriana</i>	x (29)
<i>P. snatolana</i>	x (1)
<i>P. zhuravlevii</i>	x (29)
<i>Platycarya pseudobrauni</i>	x (37)
<i>P. sp.</i>	x (29)
<i>Populus arnaudii</i>	x (29)
<i>P. budantsevii</i>	x (1, 21, 29)
<i>P. celastrophylla</i>	x (29)
<i>P. gigantophylla</i>	x (21, 22, 29)
<i>P. grandifolia</i>	x (29)
<i>P. iljinskaja</i>	x (29)
<i>P. kryshtofovichii</i>	x (29)
<i>P. microdentata</i>	x (21, 29)
<i>P. napanensis</i>	x (21)
<i>P. snatolensis</i>	x (22)
<i>P. sp.</i>	x (37)
<i>Posidonia parisiensis</i>	x (29)

<i>Pterocarya antiqua</i>	x (21)	
<i>P. pugetensis</i>	x (37)	
<i>Quercus furuhjelmi</i>	x (37)	
<i>Q. irtysheensis</i>	x (29)	
<i>Q. kornilovae</i>	x (29)	
<i>Q. stolbovii</i>	x (1)	
<i>Rhamnus</i>	x (29)	x (37)
<i>Rhododendron mainachense</i> (b)	x (1, 21)	[3]
<i>Rhus angustifolia</i>	x (29)	
<i>R. cf. mixta</i>		x (37)
<i>R. turcomanica</i>	x (25, 29)	
<i>Rubus frutalis</i> (b)	x (29)	[3]
<i>Sabal</i> (b)	x (29)	[3]
<i>Sabalites</i>		x (37)
<i>Sassafras</i> (b)	x (22, 29)	[3]
<i>Sorbaria</i>	x (29)	
<i>Spondylos</i>	x (29)	
<i>Sterculia kalkamanica</i> (b)	x (29)	[3]
<i>Ternstroemites kamtschaticus</i>	x (1)	[3]
<i>Tetracentron piperoides</i>		x (37, 39)
<i>T. sp.</i>	x (29)	
<i>Tetracera</i>	x (29)	[3]
<i>Tilia beringiana</i>	x (21)	

<i>T. carbonensis</i>	x (37)
<i>T. eojaponica</i>	x (29)
<i>T. rectinervis</i>	x (21)
<i>Trochodendroides arctica</i>	x (1, 21, 22, x (8, BC) 29)
<i>T. plutonii</i>	x (1, 21)
<i>Tsukada tiliifolia</i>	x(?)
<i>Typha</i> (b)	x (29) [3]
<i>Ulmites pacifica</i>	x (22)
<i>Ulmus compacta</i>	x (1, 21, 29)
<i>U. ezoana</i>	x (29)
<i>U. harutoriensis</i>	x (29)
<i>U. latiserrata</i>	x (1)
<i>U. longifolia</i>	x (29)
<i>U. monstrosa</i>	x (21)
<i>U.</i> sp.	x (37)
<i>Ungnandia kryshtofovichii</i>	x (29) [3]
<i>Ushia</i>	x (29)
<i>Vaccinium</i> (b)	x (29) [3]
<i>Viburnum antiquum</i> (a)	x (22)
<i>V. gracile</i>	x (21)
<i>Vitis grigorenkoi</i>	x (21)
<i>V. microdentata</i>	x (22)

<i>V. sachalinensis</i>	x (29)	
<i>V. sp.</i>		x (37)
<i>Zelkova kushiroensis</i> (b)	x (29)	[3]
<i>Z. zelkovaefolia</i>	x (29)	
<i>Zingiberites borealis</i>	x (29)	[3]
<i>Zingiberopsis</i> (b)	x (29)	
<i>Zizyphus alaskanus</i>		x (37)
<i>Z. sp.</i>	x (29)	

Oligocene

Ferns and allied groups

<i>Osmunda claytonites</i>	x (41)	
<i>O. heeri</i>	x (25)	
<i>Salvinia mildeana</i>	x (25)	[3]

Gymnosperms

<i>Cedrus</i> cf. <i>atlantica</i>	x (25)	
<i>C. perialata</i>		x (41)
<i>Glyptostrobus europaeus</i>	x (25)	
<i>G. vacuipites</i>		x (41)
<i>Metasequoia disticha</i> (5)	x (25)	
<i>M. papillapollenites</i>		x (41)
<i>Pinus flexuosa</i>	x (25)	
<i>P. sp.</i>		x (41)

<i>Sequoia abietina</i>	x (25)
<i>S. affinis</i>	x (43)
<i>Taxodium dubium</i>	x (25)
<i>T. hiatipites</i>	x (44)
<i>Taxus palaeocuspisidata</i>	x (25) [3]

Angiosperms

<i>Acer aegopodiifolium</i>	x (25)
<i>A. chaneyi</i>	x (42)
<i>A. compositifolium</i>	x (25)
<i>A. kenaicum</i>	x (42)
<i>A. kinjakense</i>	x (25)
<i>A. monoides</i>	x (25)
<i>A. paleosaccharinum</i>	x (25)
<i>A. cf. saccharinum</i>	x (25)
<i>A. subcampestre</i>	x (25)
<i>Ailanthus americana</i>	x (43)
<i>A. confucii</i>	x (25)
<i>Alnus hersutiformis</i>	x (25)
<i>A. jeroniae</i>	x (25)
<i>A. korniloviae</i>	x (25)
<i>A. aff. martini</i>	x (37)
<i>A. neuburgiae</i>	x (25)

<i>A. protomaximowiczii</i>	x (43)	
<i>A. schmalhausenii</i>	x (25)	
<i>A. verus</i>		x (41)
<i>Ampelopsis cf. delavayana</i>	x (25)	[3]
<i>A. schischkinii</i>	x (25)	
<i>Aponogeton czokusianus</i>	x (25)	
<i>A. tertiaries</i>	x (25)	
<i>Asclepias zaissanica</i>	x (25)	[3]
<i>Astronium ninae</i>	x (25)	
<i>A. truncatum</i>		x (43)
<i>Betula claripites</i>		x (41)
<i>B. subpubescens</i>	x (25)	
<i>B. thor</i>		x (43)
<i>Carex scheuchzeri</i>	x (25)	[3]
<i>Carpinus lanceolata</i> (a)	x (25)	[3]
<i>Carya juxtaporipites</i>		x (41)
<i>C. libbeyi</i>		x (43)
<i>C. sp.</i>	x (25)	
<i>Castanea antipovii</i>	x (25)	
<i>C. dolichophylla</i>		x (43)
<i>C. protocrenata</i>	x (25)	
<i>C. zaisanica</i>	x (25)	
<i>Cedrela lancifolia</i>		x (43)

<i>C. sp.</i>	x (25)
<i>Celastrus aralensis</i>	x (25)
<i>C. oregonensis</i>	x (43)
<i>Cercidiphyllum arcticum</i>	x (43)
<i>C. crenatum</i>	x (25)
<i>Coccus heteromorpha</i>	x (43)
<i>C. schischkinii</i>	x (25)
<i>Comptonia acutiloba</i>	x (25) [3]
<i>Cornus cornella</i>	x (43)
<i>C. graeffii</i>	x (25)
<i>C. studeri</i>	x (25)
<i>Corylus jarmolenkoi</i>	x (25)
<i>C. sp.</i>	x (41)
<i>Cotinus lavrovii</i> (a)	x (25)
<i>Crataegus pacifica</i>	x (43)
<i>C. sp.</i>	x (25)
<i>Cyperacites angustifolia</i>	x (25) [3]
<i>Diospyros brachysepala</i>	x (25)
<i>D. kryshtofovichii</i>	x (25)
<i>D. oregoniana</i>	x (43)
<i>Eucalyptus borissovii</i>	x (25) [3]
<i>Eucommia borissovii</i>	x (25)
<i>E. montana</i>	x (43)

<i>E. palaeoulmoides</i>	x (25)
<i>Euonymus pacificus</i>	x (43)
<i>E. proserpinae</i>	x (25)
<i>Fagus antipofii</i>	x (25)
<i>F. granulata</i>	x (41)
<i>Firmiana</i>	x (25)
<i>Fraxinus columbiana</i>	x (41)
<i>F. dubia</i>	x (25)
<i>Hemiptelea kryshtofovichii</i>	x (25)
<i>Hovenia thunbergii</i>	x (25)
<i>Juglans horniana</i>	x (41)
<i>J. sp.</i>	x (25)
<i>Laurus aff. nobilis</i>	x (25) [3]
<i>Lindera vassilenkoi (b)</i>	x (25) [3]
<i>Liquidambar europaea</i>	x (25)
<i>L. manucoica</i>	x (25)
<i>L. sp.</i>	x (41)
<i>Magnolia reticulata</i>	x (37)
<i>M. takhtajanii</i>	x (25)
<i>Mimosites acaciafolius</i>	x (43)
<i>M. sp.</i>	x (25)
<i>Myrica annulites</i>	x (41)
<i>M. cf. lignitum</i>	x (25)

<i>M. longifolia</i>	x (25)	
<i>M. serrulata</i>		x (43)
<i>Nelumbo protospeciosa</i>	x (25)	
<i>N.</i> sp.		x (43)
<i>Nyssa</i>	x (25)	x (41)
<i>Palibinia pinnatifida</i>	x (25)	
<i>Phragmites alaskana</i>		x (43)
<i>P. oenningensis</i>	x (25)	
<i>Phellodendron grandifolium</i> (b)	x (25)	
<i>Pistacia zaisanica</i>	x (25)	
<i>Platanus comstocki</i>		x (37)
<i>P. dissecta</i>		x (43)
<i>P. vassilijevii</i>	x (25)	
<i>Poacites</i>	x (25)	[3]
<i>Populus arnaudii</i>	x (25)	
<i>P. balsamoides</i>		x (43)
<i>P. mutabilis</i>	x (25)	
<i>Physocarpus</i>	x (25)	
<i>Pterocarya paradisiaca</i>	x (25)	
<i>P. stellatus</i>		x (41)
<i>Quercus alexeevii</i>	x (25)	
<i>Q. borissovii</i>	x (25)	
<i>Q. convexa</i>		x (43)

<i>Q. klinkerishica</i>	x (25)
<i>Q. oxyodon</i>	x (25)
<i>Q. palaeoserrata</i>	x (25)
<i>Q. parainfectoria</i>	x (25)
<i>Q. parazaisanica</i>	x (25)
<i>Q. patelliformis</i>	x (25)
<i>Q. protopontica</i>	x (25)
<i>Q. pseudorobur</i>	x (25)
<i>Q. shiabensis</i>	x (41)
<i>Q. zaisanica</i>	x (25)
<i>Rhamnus rectinervis</i> (a)	x (25)
<i>Rhus altynica</i>	x (25)
<i>R. kilnkerischica</i>	x (25)
<i>R. miosuccedanea</i>	x (43)
<i>R. turocomanica</i>	x (25)
<i>Salix truckeana</i>	x (43)
<i>S. sp.</i>	x (25)
<i>Sorbus praetorminalis</i> (a)	x (25) [3]
<i>Spiraea danibekii</i>	x (25) [3]
<i>Strelitzia borealis</i>	x (25)
<i>Toxicodendron junnanense</i>	x (25)
<i>Typha latissima</i>	x (25)
<i>T. lesquereuxi</i>	x (43)

<i>Ulmus carpinoides</i>	x (25)	
<i>U. drepanodonta</i>	x (25)	
<i>U. paucidentata</i>		x (43)
<i>U. schischkinii</i>	x (25)	
<i>Vitis</i> cf. <i>vulpina</i> (a)	x (25)	[3]
<i>Zelkova ungeri</i>		x (43)
<i>Z. zelkovifolia</i>	x (25)	

Miocene

Ferns and allied groups

<i>Dryopteris</i>	x (30)	x (35, 45)
<i>Equisetum</i>	x (47)	x (35)
<i>Onoclea sensibilis</i>	x (47)	x (35, 45)
<i>Osmunda doroshiana</i>	x (47)	
<i>O. heeri</i>	x (25)	
<i>O.</i> sp.		x (44)
<i>Salvinia mildeana</i>	x (25)	[3]

Gymnosperms

<i>Ginkgo biloba</i> (7)		x (35, 45)
<i>Glyptostrobus europaeus</i>	x (25, 47)	x (35, 45)
<i>Keteleeria</i>	x (30)	x (46)
<i>Larix</i> cf. <i>L. omoloica</i>	x (47)	

<i>L.</i> sp.	x (30)	x (44)
<i>Metasequoia disticha</i>	x (25, 47)	
<i>M. glyptostroboides</i>		x (35, 45)
<i>Picea koefiensis</i>	x (47)	
<i>P.</i> sp.		x (44)
<i>Pinus</i> cf. <i>ponderosa</i>		x (35)
<i>P.</i> sp.	x (30)	
<i>Pseudolarix fossilis</i>	x (25)	
<i>P.</i> sp.	x (47)	x (36)
<i>Sequoia sempervirens</i> (8)		x (35)
<i>S.</i> sp.		x(44)
<i>Taxodium distichum</i>		x (35)
<i>T. dubium</i>	x (25, 47)	

Angiosperms

<i>Acer aegopodiifolium</i>	x (25)
<i>A. chaneyi</i>	x (42)
<i>A. compositifolium</i>	x (25)
<i>A. ezoanum</i>	x (35)
<i>A. monoides</i>	x (25)
<i>A. palaeosaccharinum</i>	x (25)
<i>Alangium mikii</i>	x (45)
<i>A.</i> sp.	x (30)

<i>Alfaropsis</i>	x (30)	
<i>Alnus barnesi</i>	x (35)	
<i>A. feroniae</i>	x (25)	
<i>A. hirsutiformis</i>	x (25)	
<i>A. kerniloviae</i>	x (25)	
<i>A. schmalhausenii</i>	x (25)	
<i>Apocynophyllum helveticum</i>	x (25)	[3]
<i>Aponogeton czokusianus</i>	x (25)	
<i>Betula sublutea</i>	x (35)	
<i>B. subpubescens</i>	x (25)	
<i>Carex</i>	x (28)	[3]
<i>Carpinus cappensis</i>	x (35)	
<i>C. sp.</i>	x (30)	
<i>Carya bendieri</i>	x (35)	
<i>C. sp.</i>	x (30)	
<i>Castanea antipovii</i> (b)	x (25)	[3]
<i>Celastrus aralensis</i> (a)	x (25)	[3]
<i>Celtis japeti</i>	x (25)	
<i>C. sp.</i>	x (44, 45, 46)	
<i>Cercidiphyllum crenatum</i>	x (25, 47)	x (35)
<i>Cercis turgaica</i>	x (25)	[3]
<i>Cissus</i> (b)	x (30)	[3]
<i>Cladrastis japonica</i>	x (35, 46)	

<i>C. sp.</i>	x (30)	
<i>Comptonia acutiloba</i>	x (25)	
<i>C. cf. naumani</i>		x (35)
<i>Cornus graeffii</i>	x (25)	
<i>C. sp.</i>		x (35)
<i>Corylus harrimani</i>		x (35)
<i>C. jarmolenkoi</i>	x (25)	
<i>Cotinus lavrovii</i>	x (25)	
<i>Crataegus itelmenensis</i>	x (47)	
<i>C. sp.</i>	x (25)	x (35, 46)
<i>Davidia</i>	x (30)	
<i>Desmodium</i>	x (30)	[3]
<i>Eucommia cf. montana</i>		x (45)
<i>E. sp.</i>	x (30)	
<i>Fagus antipofii</i>	x (25)	x (35)
<i>F. chankaica</i>	x (30)	
<i>F. evenensis</i>	x (30)	
<i>Fraxinus dubia</i>	x (25)	
<i>F. kenaica</i>		x (45)
<i>Hovenia thunbergii</i>	x (25)	
<i>Hydrangea minima</i>	x (47)	
<i>H. sp.</i>		x (46)
<i>Idesia</i>	x (30)	[3]

<i>Ilex</i>	x (30)	x (44)
<i>Indigofera</i>	x (30)	
<i>Juglans</i>	x (25, 30)	
<i>J. type (9)</i>		x (44)
<i>Liquidambar europaea</i>	x (25)	
<i>L. mioformosana</i>		x (35)
<i>Lindera</i>	x (30)	[3]
<i>Litsea</i>	x (30)	[3]
<i>Magnolia takhtajanii</i>	x (25)	[3]
<i>Meliosma</i>	x (30)	[3]
<i>Microtropis</i>	x (30)	
<i>Myrica czokusiana</i>	x (25)	
<i>M. sp.</i>		x (35, 46)
<i>Nelumbo protospeciosa</i>	x (25)	[3]
<i>Nuphar</i>	x (25)	x (35, 46)
<i>Nyssa knowltoni</i>		x (35)
<i>N. sp.</i>	x (30)	x (35, 44, 46)
<i>Ostrya</i>	x (30)	[3]
<i>Paliurus</i>	x (30)	[3]
<i>Parrotia prisitina</i>	x (25)	[3]
<i>Periploca kryshtofocichii</i>	x (25)	
<i>Phellodendron grandifolium</i>	x (25, 47)	
<i>Phragmites oeningensis</i>	x (25)	[3]

<i>Platanus bendirei</i> (a)		x (35, 45)
<i>Populus kenaiana</i>		x (35)
<i>P. populina</i>	x (25)	
<i>Potamogeton alaskanus</i>		x (45)
<i>P.</i> sp.	x (28, 44)	x (35, 46)
<i>Prunus rubeshibensis</i>	x (47)	
<i>P.</i> sp.		x (46)
<i>Pterocarpa mixta</i>		x (35)
<i>P. paradisiaca</i>	x (25)	
<i>Pterocarya nigella</i>		x (44)
<i>P. kamtschatica</i>	x (47)	
<i>Quercus alexeevii</i>	x (25)	
<i>Q. bretzi</i>		x (35)
<i>Q. pseudorobur</i>	x (25)	
<i>Rhamnus</i>		x (46)
<i>R.</i> sp.	x (47)	
<i>Rhus altynica</i>	x (25)	
<i>R. aff. glabra</i>		x (35)
<i>Rosa</i>	x (47)	x (46)
<i>Salix cappensis</i>		x (35)
<i>S. itelmenensis</i>	x (47)	
<i>Scirpus</i>	x (25)	[3]
<i>Sorbus gabrieljaniae</i>	x (25)	[3]

<i>Sparganium</i>	x (28)	x (44, 46)
<i>Spatholobus</i>	x (30)	
<i>Spiraea korfiensis</i>	x (47)	
<i>S. sp.</i>		x (46)
<i>Strelitzia borealis</i>	x (25)	
<i>Tilia</i>	x (25)	x (35, 44, 46)
<i>Trapa</i>	x (25)	x (46)
<i>Trochodendron</i>	x(38, 39)	x (38,39)
<i>Typha latissima</i>	x (25)	
<i>T. sp.</i>		x (35, 44, 46)
<i>Ulmus carpinoides</i>	x (25)	
<i>U. drepanodonta</i>	x (25)	
<i>U. longifolia</i>		x (35, 46)
<i>Zelkova oregoniana</i>		x (35)
<i>Z. sp.</i>	x (30)	

Pliocene

Ferns and allied groups

<i>Lycopodium</i>	(w)	x (48)
<i>Pteridium</i>	(w)	x (48)
<i>Selaginella</i>	(w)	x (48)

Gymnosperms

<i>Abies</i>	x (26, 28)	x (48)
<i>Ephedra</i>	x (27)	

<i>Larix</i>	x (26, 29)	x (48)
<i>Picea</i>	x (26, 28, 29)	x (48)
<i>Pinus</i>	x (26, 28, 29)	x (48)
<i>Taxodium</i>	x (26)	
<i>Tsuga</i>	x (26, 28, 29)	

Angiosperms

<i>Acer</i>	x (26)	[3]
<i>Alnaster</i>	x (29)	
<i>Alnus</i>	x (26, 28, 29)	[3]
<i>Anromeda</i>	x (27)	x (48)
<i>Aralia</i>	x (26, 27, 28)	[3]
<i>Artemisia</i>	x (26, 27)	(w)
Asteraceae	x (26)	
<i>Betula</i> sect. <i>alba</i>	x (26)	
<i>B. humulis</i>	x (26)	
<i>Betula</i>		x (48)
<i>Bunias cochlearrioides</i>	x (26)	
<i>Carpinus</i>	x (27, 28, 29)	[3]
<i>Carya</i>	x (26)	[3]
Chenopodiaceae	x (26, 27)	(w)
<i>Comptonia</i>	x (27)	[3]
<i>Corylus</i>	x (27, 29)	x (48)

<i>Diervilla</i>	x (27)	
Ericaceae	x (28, 29)	x (48)
<i>Fagus</i>	x (26, 28)	[3]
<i>Fraxinus</i> (a)		x (48)
<i>Ilex</i>	x (26, 29)	x (48)
<i>Juglans</i>	x (29)	[3]
<i>Lemna</i>		x (48)
<i>Liquidambar</i>	x (26)	[3]
<i>Magnolia</i>	x (26)	[3]
<i>Morus tanaitica</i>	x (26, 27)	[3]
<i>Myrica</i>	x (28, 29)	[3]
<i>Myriophyllum</i>		x (48)
<i>Nyssa</i>	x (26)	[3]
<i>Ostrya</i>	[52]	x (48)
<i>Parrotia</i>	x (26)	[3]
<i>Pittosporum</i>	x (26)	
Plumbaginaceae	x (27)	
Poaceae	x (27, 28)	x (48)
<i>Polycnemum</i>	x (26)	
<i>Polygonum</i>		x (48)
<i>Pterocarya</i>	x (26)	[3]
<i>Quercus</i>	x (26, 27)	x (48)
<i>Rubus</i>	x (26)	[3]

<i>Rumex marshallianus</i>	x (26)	[3]
<i>Salix</i>	x (26)	[3]
<i>Sambucus</i>	x (27)	[3]
<i>Saxifraga</i>		x (48)
<i>Staphylea</i>	x (26)	[3]
<i>Tilia</i>	x (26, 27, 29)	[3]
<i>Typha</i> (a)		x (48)
<i>Ulmus</i>	x (26, 27)	[3]
<i>Urtica</i>	[52]	x (48)
<i>Vitis</i>	x (27)	[3]

Notes to Appendix 2.

1. In Zhilin (1991, ed., p. 77, table 3) ages in the three columns translate as early Oligocene, late Oligocene, and early Miocene. In table 4 (p. 77) the ages are late Eocene, early Oligocene, late Oligocene, and early Miocene. Table 4 (p. 119) is a comparison of early Miocene vegetation at three different localities.
2. *Protophyllum* and *Aspidiophyllum* now transferred to *Pseudoprotophyllum* fide Vakhrameed (1991, p. 182).
3. See Rothwell and Stockey (1994) for report of *H. pinnata* from the Cretaceous St. Mary River Formation, southern Alberta, Canada.
4. See Pigg et al. (2003) for reports of *Carpinus*, *Corylus*, and *Palaeocarpinus* from the Eocene of northwestern North America.

5. See LePage et al. (2005) for a summary of the fossil record and ecology of *Metasequoia*. Eocene occurrences in the Canadian High Arctic (Axel Heiberg Island) and in the early Oligocene Bridge Creek flora of Oregon are discussed by Richter and LePage and by Meyer in LePage et al. (2005).
6. *Pinus* was thought to have originated in the Cretaceous (Albian, ca. 100 Ma) and spread from centers in eastern North America and Western Europe (Millar, 1999). Its presence in Beringia at 133-140 Ma pre-dates this time (Falcon-Lang et al., 2016).
7. The oldest *Ginkgo* (as opposed to Ginkgophytes) is from the Lower Jurassic (ca. 190 Ma) in the Asiatic part of the former USSR (<http://www.ucmp.berkeley.edu/seedplants/ginkgoales/ginkgofr.html>; Taylor, 2009, chapter 18).
8. The oldest *Sequoia* fide Ahuja and Neale (2002) is from the Jurassic of south Manchuria.
9. *Juglans* as a genus dates from the middle Eocene (ca. 45 Ma) of North America (Stanford et al., 2000; see also Aradhya, 2006; Manchester, 1987; Wilson et al.-
<http://people.brandon.ca/greenwood/files/2013/wilsonbasinger-small.pdf>).).
10. The North American occurrences listed in [3] is ‘Paleocene or Eocene’, Kodiak Island, Alaska (Hollick, 1936).
11. The North American occurrences listed in [3] are Paleocene, Spitzbergen (Heer, 1870, 1883).
12. See Dilcher (1969), Eocene, southeastern U.S.; Graham and Jarzen (1969, Oligocene, Puerto Rico; Graham, 1976, Veracruz, Mexico; Graham et seq.).
13. Reported from the Cretaceous (Campanian, ca. 83-70 Ma), Vancouver Island, Canada, Stockey and Rothwell (2003).
14. Earliest reported occurrence of *Ilex* as a genus is in the Paleocene (65 Ma) of Australia but placement is uncertain and Beaulieu et al. (2013) accept it as evidence of the family

Aquifoliaceae with the broader Campanulids originating in the middle Albian (Cretaceous, ca. 105 Ma) in the Southern Hemisphere (see also Manen et al., 2002).

- a. Indicates listed elsewhere in the Appendix in both eastern and western Beringia.
- b. Indicates oldest occurrence for the genus as listed in the Appendix is western Beringia and vicinity.
- c. Op cit.- eastern Beringia and vicinity.

References to Appendix 2

- Ahuja, M. R. and D. B. Neale. 2002. Origins of polyploidy in coast redwood (*Sequoia sempervirens*) and relationship of coast redwood to other genera of Taxodiaceae. *Silvae Genetica* 51: 93-100.
- Akhmetiev, M. A. 2007. Paleocene and Eocene floras of Russia and adjacent regions: climatic conditions of their development. *Paleontological Journal* 41: 1032-1039.
- 2010. Paleocene and Eocene floristic and climatic change in Russia and northern Kazakhstan. *Bulletin of Geosciences* 85: 77-94.
- Aradhya, M. K., D. Potter, and C. J. Simon. 2006. Origin, evolution, and biogeography of *Juglans*: a phylogenetic perspective. In: M. E. Malvolti and D. Avanzato (eds.), *Proceedings of the Vth International Walnut Symposium. Acta Horticulturae* 705: 85-94.
- Arkhipov, S. A., V. S. Volkova, I. D. Zolnikov, V. S. Zykina, A. A. Krukover, and L. A. Kul'kova. 2005. West Siberia. In: A. A. Velichko and V. P. Nechaev (eds.). 2005. *Cenozoic Climatic and Environmental Changes in Russia*. Geological Society of America Special Paper 382. Pp. 67-88.
- Beaulieu, J. M., D. C. Tank, and M. J. Donoghue. 2013. A Southern Hemisphere origin for campanulid angiosperms, with traces of the break-up of Gondwana. *BMC Evolutionary Biology* 13: 80.
- Becker, H. F. 1969. Fossil plants of the Tertiary Beaverhead Basins in southwestern Montana. *Palaeontographica* 127, Abt. B, 1: 1-142.
- Bell, W. A. 1949. Uppermost Cretaceous and Paleocene floras of western Alberta. Canada Department of Mines and Resources, Geological Survey Bulletin No. 13.

Ottawa.

- Berry, E. W. 1926. Tertiary floras from British Columbia. Canada Department of Mines, Geological Survey Bulletin 42: 91-116.
- Boyd, A. 1990. *Musopsis*, a banana-like leaf genus from the Early Tertiary of eastern North Greenland. American Journal of Botany 79: 1359-1367.
- Brink, K. S., R. A. Stockey, G. Beard, and W. C. Wehr. 2009. *Cunninghamia hornbyensis* sp. nov.: permineralized twigs and leaves from the Upper Cretaceous of Hornby Island, British Columbia, Canada. Review of Palaeobotany and Palynology 155: 89-98.
- Budantsev, L. Y. (ed.). 1994. Magnoliophyta Fossilia Rossiae Civitatum Finitimarum. Vol. 3. Leitneriaceae-Juglandaceae. Institutum Botanicum Komarovianum, Academiae Scientiarum Rossicum, St. Petersburg.
- 1997. Late Eocene Flora of Western Kamchatka. Russian Academy of Sciences, Proceedings of Komarov Botanical Institute, Issue 19, St. Petersburg.
- 2005. Magnoliophyta Fossilia Rossiae Civitatum Finitimarum. Vol. 4. Nyctaginaceae-Salicaceae. Institutum Botanicum Komarovianum, Academiae Scientiarum Rossicum, St. Petersburg.
- Chaney, R. W. 1951. A revision of fossil *Sequoia* and *Taxodium* in western North America based on the recent discovery of *Metasequoia*. Transactions of the American Philosophical Society 40: 171-235.
- Chelebaeva, A. I. 1978. Miocene floras of Vostok Kamchatka. Nauka, Moscow. 154 pp. [in Russian].
- Dilcher, D. L. 1969. *Podocarpus* from the Eocene of North America. Science 164: 299-

301.

Doyle, J. A. and L. J. Hickey. 1976. Pollen and leaves from the mid-Cretaceous Potomac Group and their bearing on early angiosperm evolution. In: C. B. Beck (ed.), Origin and Early Evolution of Angiosperms. Columbia University Press, New York. Pp. 139-206.

Falcon-Lang, H. J., V. Mages, and M. Collinson. 2016. The oldest *Pinus* and its preservation by fire. *Geology* 44: 303-306.

Fradkina, A. F., M. N. Alekseev, A. A. Andreev, and V. A. Klimanov. 2005a. East Siberia (based on data obtained mainly in Central Yakutia). In: A. A. Velichko and V. P. Nechaev (eds.). *Cenozoic Climatic and Environmental Changes in Russia*. Geological Society of America Special Paper 382. Pp. 89-103.

-----, O. V. Grinenko, S. A. Laukhin, V. P. Nechaev, A. A. Andreev, and V. A. Klimanov. 2005b. Northeastern Asia. In: A. A. Velichko and V. P. Nechaev (eds.). *Cenozoic Climatic and Environmental Changes in Russia*. Geological Society of America Special Paper 382. Pp. 105-120.

Golovneva, L. B. 2000. Aquatic plant communities at the Cretaceous-Paleogene boundary in north-eastern Russia. *Acta Palaeobotanica* 40: 139-151.

Graham, A. 1976. Studies in neotropical paleobotany. II. The Miocene communities of Veracruz, Mexico. *Annals of the Missouri Botanical Garden* 63: 787-842.

-----, 2010. Late Cretaceous and Cenozoic History of Latin American Vegetation and Terrestrial Environments. Missouri Botanical Garden Press, St. Louis.

-----, 2012. Catalog and literature guide for Cretaceous and Cenozoic vascular plants of the New World. *Annals of the Missouri Botanical Garden* 98: 539-541.

- and D. M. Jarzen. 1969. Studies in neotropical paleobotany. I. The Oligocene communities of Puerto Rico. Annals of the Missouri Botanical Garden 56: 308-357.
- Harris, A. J., M. Papes, Y.-D. Gao, and L. Watson. 2014. Estimating paleoenvironments using ecological niche models of nearest living relatives: a case study of Eocene *Aesculus* L. Journal of Systematics and Evolution 52: 16-34.
- Heer, O. 1870. Flora fossilis Arctica. Vol 2, pt. 2. J. Wurster, Zurich.
- 1855. Flora Tertiaria Helvetiae, bd. 1. Winterthur.
- 1876. Beitrage zur fossilen flora Spitzbergens. Kunglia Svenska Vetenskaps-Akademiens Handlingar 14: 3-93.
- 1883. Flora Fossilis Groenlandica. Pt. 2, No. 4-6, Flora Fossilis Arctica, Vol. 7, pp. 148-275. Zurich.
- Herman, A. B. 1990. Late Cretaceous floras and climate of the Anadyr-Koryakian subregion (north-east USSR). Proceedings of the Symposium Paleofloristic and Paleoclimatic Changes in the Cretaceous and Tertiary, International Geological Correlation Programme, Project No. 216, Global Biological Events in Earth History, Prague, 1989. Pp. 73-79.
- 2013. Albian-Paleocene flora of the North Pacific: systematic composition, palaeofloristics and phytostratigraphy. Stratigraphy and Geological Correlation 21: 689-747.
- , M. G. Moiseeva, R. A. Spicer, and A. Ahlberg. 2004. Maastrichtian-Paleocene floras of northeastern Russia and north Alaska and floral changes at the Cretaceous-Paleogene boundary. Stratigraphy and Geological Correlation 12: 485-494.
- and R. A. Spicer. 2010. Mid-Cretaceous floras and climate of the Russian high

- Arctic (Novosibirsk Islands, northern Yakutiya). Palaeogeography, Palaeoclimatology, Palaeoecology 295: 409-422.
- Hollick, A. 1930. The Upper Cretaceous floras of Alaska. U.S. Geological Survey Professional Paper 159. 116 pp.
- 1936. The Tertiary floras of Alaska. U.S. Geological Survey Professional Paper 182. 185 pp.
- Knowlton, F. H. 1899. Fossil flora of Yellowstone National Park. U.S. Geological Survey Monograph 32: 651-882.
- 1910. Descriptions of fossil plants from the Mesozoic and Cenozoic of North America. I. Smithsonian Miscellaneous Collection 52: 489-496.
- LaMotte, R. S. 1952. Catalogue of the Cenozoic Plants of North America through 1950. Geological Society of America Memoir 51, New York.
- Lavrenko, O. D. and L. I. Fot'janova. 1994. Some early Paleogene species from western Kamchatka. In: M. C. Boulter and H. C. Fisher (eds.), Cenozoic Plants and Climates of the Arctic. NATO ASI Series, Series 1: Global Environmental Change, Vol. 27. Pp. 315-321.
- Lebedev, Ye. L. 1992. The Cretaceous floras of northeastern Asia. International Geology Review 34: 794-805.
- Leopold, E. B. and G. Liu. 1994. A long pollen sequence of Neogene age, Alaska Range. Quaternary International 22/23: 103-140.
- LePage, B. A., C. J. Williams, and H. Yang (eds.). 2005. The Geobiology and Ecology of *Metasequoia*. Springer, Berlin.
- Lesquereux, L. 1874. Contribution to the fossil flora of the Western Territories, pt. 1.

- U.S. Geological Survey Report 6, Washington, D.C. 136 pp.
- 1878. Contributions to the fossil flora of the Western Territories, pt. 2. U.S. Geological Survey Report 7, Washington, D.C. 366 pp.
- Manchester, S. R. 1987. The fossil history of the Juglandaceae. Monographs in Systematic Botany from the Missouri Botanical Garden 21: 1-137.
- , Q.-Y. Xiang, T. M. Kodrul, and M. A. Akhmetiev. 2009. Leaves of *Cornus* (Cornaceae) from the Paleocene of North America and Asia confirmed by trichome characters. International Journal of Plant Sciences 170: 132-142.
- Manen, J.-F., M. C. Boulter, and Y. Naciri-Graven. 2002. The complex history of the genus *Ilex* L. (Aquifoliaceae): evidence from the comparison of plastid and nuclear DNA sequences and from fossil data. Plant Systematics and Evolution 235: 79-98.
- McIver, E. E. and J. F. Basinger. 1993. Flora of the Ravenscrag Formation (Paleocene), southwestern Saskatchewan, Canada. Palaeontographica Canadiana 10: 1-167 pp.
- Meyer, H. W. 2005. *Metasequoia* in the Oligocene Bridge Creek flora of western North America: ecological implications and the history of research. In: B. A. LePage, C. J. Williams, and H. Yang (eds.). The Geobiology and Ecology of *Metasequoia*. Springer, Berlin. Pp. 159-186.
- Millar, C. I. 1999. Evolution and biogeography of *Pinus radiata*, with a proposed revision of its Quaternary history. New Zealand Journal of Forestry Science 29: 335-365.
- Moiseeva, M. G., A. B. Herman, and R. A. Spicer. 2009. Late Paleocene flora of the northern Alaska Peninsula. The role of Transberingian plant migrations and climate change. Paleontological Journal 43: 1298-1308.

- Pavlyutkin, B. I. and I. Yu. Chekryzhov. 2007. Correlation of Miocene floras of the Climatic Optimum in the Far East. *Doklady Earth Sciences* 417: 1156-1159.
- Penhallow, D. P. 1908. Report on Tertiary plants of British Columbia. Canada Department of Mines, Geological Survey Branch, No. 1013, Ottawa.
- Piel, K. M. 1971. Palynology of Oligocene sediments from central British Columbia. *Canadian Journal of Botany* 49: 1885-1920.
- Pigg, K. B., R.M. Dillhoff, M. L. DeVore, and W. C. Wehr. 2007. New diversity among Trochodendraceae from the early/middle Eocene Okanogan Highlands of British Columbia, Canada, and northeastern Washington State, United States. *International Journal of Plant Sciences* 168: 521-532.
- , S. R. Manchester, and W. C. Wehr. 2003. *Corylus*, *Carpinus*, and *Palaeocarpinus* (Betulaceae) from the middle Eocene Klondike Mountain and Allenby formations of northwestern North America. *International Journal of Plant Sciences* 162: 1187-1198.
- , W. C. Wehr, and S. M. Ickert-Bond. 2001. *Trochodendron* and *Nordenskioldia* (Trochodendraceae) from the middle Eocene of Washington State, USA. *International Journal of Plant Sciences* 162: 1187-1198.
- Pound, M. J., R. L. Lowther, J. Peakall, R. J. Chapman, and U. Salzmann. 2015. Palynological evidence for a warmer boreal climate in the late Pliocene of the Yukon Territory, Canada. *Palynology* 39: 91-102.
- Reinink-Smith, L. M. and E. B. Leopold. 2005. Warm climate in the late Miocene of the south coast of Alaska and the occurrence of Podocarpaceae pollen. *Palynology* 29: 205-262.
- Retallick, G. J. and D. L. Dilcher. 1981. A coastal hypothesis for the dispersal and rise

- to dominance of flowering plants. In K. J. Niklas (ed.), Paleobotany, Paleoecology and Evolution, Vol. 2. Praeger, New York. Pp. 27-77.
- Richter, S. L. and B. A. LePage. 2005. A high-resolution palynological analysis, Axel Heiberg Island, Canadian High Arctic. In: B. A. LePage, C. J. Williams, and H. Yang (eds.), The Geobiology and Ecology of *Metasequoia*. Springer, Berlin. Pp. 137-158.
- Rothwell, G. W. and R. A. Stockey. 1994. The role of *Hydropteris pinnata* gen et sp. nov. in reconstructing the cladistics of heterosporous ferns. American Journal of Botany 81:479-492.
- Shezepetov, S. V. and L. B. Golovneva. 2010. The genus *Heilungia* Prynada (Cacadopsida) in the Late Cretaceous floras of the Okhotsk-Chukotka volcanic belt (northeastern Russia). Paleontological Journal 44: 1339-1347.
- Spicer, R. A., K. S. Davies, and A. B. Herman. 1994. Circum-Arctic plant fossils and the Cretaceous-Tertiary transition. In: M. C. Boulter and H. C. Fisher (eds.), Cenozoic Plants and Climates of the Arctic. NATO ASI Series, Series 1: Global Environmental Change, Vol. 27. Pp. 161-169.
- and A. B. Herman. 1996. *Nilssoniocladus* in the Cretaceous Arctic: new species and biological insights. Review of Palaeobotany and Palynology 92: 229-243.
- and ----- . 2001. The Albian-Cenomanian flora of the Kukpowruk River, western North Slope, Alaska: stratigraphy, palaeofloristics, and plant communities. Cretaceous Research 22: 1-40.
- Stanford, A. M., R. Harden, and C. R. Parks. 2000. Phylogeny and biogeography of *Juglans* (Juglandaceae) based on *matK* and ITS sequence data. American Journal of Botany 87: 872-882.

- Stockey, R. A., G. L. Hoffman, and G. W. Rothwell. 1997. The fossil monocot *Limnobiophyllum scutatum*: resolving the phylogeny of Lemnaceae. American Journal of Botany 84: 355-368.
- and G. W. Rothwell. 2003. Anatomically preserved *Williamsonia* (Williamsoniaceae): evidence for Bennettitalean reproduction in the Late Cretaceous of western North America. International Journal of Plant sciences 164: 251-262.
- , -----, and A. B. Falder. 2001. Diversity among taxodioid conifers: *Metasequoia foxii* sp. nov. from the Paleocene of central Alberta, Canada. International Journal of Plant Sciences 162: 221-234.
- Takhtajan, A. (ed.). 1974. Magnoliophyta Fossilia URSS. Vol. 1. Magnoliaceae-Eucommiaceae. Nauka, Leningrad.
- 1982. Magnoliophyta URSS. Vol. 2. Ulmaceae-Betulaceae. Nauka, Leningrad.
- Taylor, T. N., E. L. Taylor, and M. Krings. 2009. Paleobotany, the Biology and Evolution of Fossil Plants. Academic Press, New York.
- Vakhrameev, V. A. 1958. Regional stratigraphy of the USSR. Vol. 3. Stratigraphy and fossil flora from the Jurassic and Cretaceous deposits of Velui Depression and adjacent parts of Verkhoyansk Marginal Trough. USSR Academy of Science, Moscow.
- 1991. Jurassic and Cretaceous Floras and Climates of the Earth. Cambridge University Press, Cambridge (English translation of 1988 Russian edition).
- Velichko, A. A. et al. (+ 15 authors). 2005. East European Plain. In: A. A. Velichko and V. P. Nechaev (eds.), Cenozoic Climatic and Environmental Changes in Russia. Geological Society of America Special paper 382. Pp. 31-66.
- Wolfe, J. A. 1966. Tertiary plants from the Cook Inlet region, Alaska. U.S. Geological

- Survey Professional Paper 398-B. 32 pp. [see also Wolfe et al., 1966].
- 1977. Paleogene floras from the Gulf of Alaska Region. U.S. Geological Survey Professional Paper 997. 108 pp.
- , D. M. Hopkins, and E. B. Leopold. 1966. Tertiary stratigraphy and paleobotany of the Cook Inlet Region, Alaska. U.S. Geological Survey Professional Paper 398-A. 29 pp.
- and T. Tanai. 1980. The Miocene Seldovia Point flora from the Kenai Group, Alaska. U.S. Geological Survey Professional Paper 1105. 52 pp.
- and -----. 1987. Systematics, phylogeny, and distribution of *Acer* (maples) in the Cenozoic of western North America. Journal of the Faculty of Sciences, Hokkaido University, Series 4, Geology and Mineralogy 22: 1-246.
- and W. Wehr. 1987. Middle Eocene dicotyledonous plants from Republic, northeastern Washington. U.S. Geological Survey Bulletin 1597. 25 pp.
- Zhilin, S. G. (ed.). 1991. Development of the Flora in Kazakhstan and Russian Plain from the Eocene to the Miocene. Kryshtofovich Lecture Series No. 2. Academy of Sciences of the USSR/Komarov Botanical Institute. Leningrad.